

## WO2005028469

Publication Title:

QUINAZOLINE DERIVATIVES

Abstract:

Abstract of WO2005028469

The invention concerns quinazoline derivatives of Formula (I): wherein each of R<1>, X<1>, R<2>, R<3>, R<5>, n and m have any of the meanings defined in the description; processes for their preparation, pharmaceutical compositions containing them and their use in the manufacture of a medicament for use as an antiproliferative agent in the prevention or treatment of tumours which are sensitive to inhibition of EGF and erbB receptor tyrosine kinases. Data supplied from the esp@cenet database - Worldwide

-----  
Courtesy of <http://v3.espacenet.com>

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
31 March 2005 (31.03.2005)

PCT

(10) International Publication Number  
**WO 2005/028469 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 401/12**,  
A61K 31/517, A61P 35/00

AstraZeneca Pharma, Z.I. la Pompelle, BP 1050, F-51689  
Reims (FR). **BARLAAM, Bernard, Christophe** [FR/FR];  
AstraZeneca Pharma, Z. I. la Pompelle BP 1050, Cedex  
02, F-51689 Reims (FR).

(21) International Application Number:  
PCT/GB2004/003937

(74) Agent: **ASTRAZENECA**; Global Intellectual Property,  
S-SE-151 85 Sodertalje (SE).

(22) International Filing Date:  
15 September 2004 (15.09.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
03292309.6 19 September 2003 (19.09.2003) EP  
04291248.5 14 May 2004 (14.05.2004) EP

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,  
PH, PI, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.

(71) Applicant (for AE, AG, AL, AM, AT, AU, AZ, BA, BB, BE,  
BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CY, CZ,  
DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MK, MN, MW, MX, MZ,  
NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD,  
SE, SG, SK, SL, SY, SZ, TJ, TM, TN, TR, TT, TZ, UA, UG,  
UZ, VC, VN, YU, ZA, ZM, ZW only): **ASTRAZENECA**  
AB [SE/SE]; Sodertalje, S-SE-151 85 (SE).

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,  
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,  
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for MG only): **ASTRAZENECA UK LIM-  
ITED** [GB/GB]; 15 Stanhope Gate, London Greater Lon-  
don W1K 1LN (GB).

**Published:**


- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BRADBURY,**  
**Robert, Hugh** [GB/GB]; AstraZeneca R % D Alderley,  
Alderley Park, Macclesfield Cheshire SK10 4TG (GB).  
**HENNEQUIN, Laurent, Francois, Andre** [FR/FR];

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: QUINAZOLINE DERIVATIVES



(I)

(57) Abstract: The invention concerns quinazoline derivatives of Formula (I): wherein each of R<sup>1</sup>, X<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, n and m have any of the meanings defined in the description; processes for their preparation, pharmaceutical compositions containing them and their use in the manufacture of a medicament for use as an antiproliferative agent in the prevention or treatment of tumours which are sensitive to inhibition of EGF and erbB receptor tyrosine kinases.

### QUINAZOLINE DERIVATIVES

The invention concerns certain novel quinazoline derivatives, or pharmaceutically acceptable salts thereof, which possess anti-tumour activity and are accordingly useful in methods of treatment of the human or animal body. The invention also concerns processes for the manufacture of said quinazoline derivatives, to pharmaceutical compositions containing them and to their use in therapeutic methods, for example in the manufacture of medicaments for use in the prevention or treatment of solid tumour disease in a warm-blooded animal such as man.

10 Many of the current treatment regimes for diseases resulting from the abnormal regulation of cellular proliferation such as psoriasis and cancer, utilise compounds that inhibit DNA synthesis and cellular proliferation. To date, compounds used in such treatments are generally toxic to cells however their enhanced effects on rapidly dividing cells such as tumour cells can be beneficial. Alternative approaches to these cytotoxic anti-tumour agents  
15 are currently being developed, for example selective inhibitors of cell signalling pathways. These types of inhibitors are likely to have the potential to display an enhanced selectivity of action against tumour cells and so are likely to reduce the probability of the therapy possessing unwanted side effects.

Eukaryotic cells are continually responding to many diverse extracellular signals that  
20 enable communication between cells within an organism. These signals regulate a wide variety of physical responses in the cell including proliferation, differentiation, apoptosis and motility. The extracellular signals take the form of a diverse variety of soluble factors including growth factors as well as paracrine and endocrine factors. By binding to specific transmembrane receptors, these ligands integrate the extracellular signal to the intracellular  
25 signalling pathways, therefore transducing the signal across the plasma membrane and allowing the individual cell to respond to its extracellular signals. Many of these signal transduction processes utilise the reversible process of the phosphorylation of proteins that are involved in the promotion of these diverse cellular responses. The phosphorylation status of target proteins is regulated by specific kinases and phosphatases that are responsible for the  
30 regulation of about one third of all proteins encoded by the mammalian genome. As phosphorylation is such an important regulatory mechanism in the signal transduction process, it is therefore not surprising that aberrations in these intracellular pathways result in

- 2 -

abnormal cell growth and differentiation and so promote cellular transformation (reviewed in Cohen *et al.*, Curr Opin Chem Biol, 1999, 3, 459-465).

It has been widely shown that a number of these tyrosine kinases are mutated to constitutively active forms and/or when over-expressed result in the transformation of a variety of human cells. These mutated and over-expressed forms of the kinase are present in a large proportion of human tumours (reviewed in Kolibaba *et al.*, Biochimica et Biophysica Acta, 1997, 133, F217-F248). As tyrosine kinases play fundamental roles in the proliferation and differentiation of a variety of tissues, much focus has centred on these enzymes in the development of novel anti-cancer therapies. This family of enzymes is divided into two groups - receptor and non-receptor tyrosine kinases e.g. EGF Receptors and the SRC family respectively. From the results of a large number of studies including the Human Genome Project, about 90 tyrosine kinase have been identified in the human genome, of this 58 are of the receptor type and 32 are of the non-receptor type. These can be compartmentalised in to 20 receptor tyrosine kinase and 10 non-receptor tyrosine kinase sub-families (Robinson *et al.*, Oncogene, 2000, 19, 5548-5557).

The receptor tyrosine kinases are of particular importance in the transmission of mitogenic signals that initiate cellular replication. These large glycoproteins, which span the plasma membrane of the cell possess an extracellular binding domain for their specific ligands (such as Epidermal Growth Factor (EGF) for the EGF Receptor). Binding of ligand results in the activation of the receptor's kinase enzymatic activity that is encoded by the intracellular portion of the receptor. This activity phosphorylates key tyrosine amino acids in target proteins, resulting in the transduction of proliferative signals across the plasma membrane of the cell.

It is known that the erbB family of receptor tyrosine kinases, which include EGFR, erbB2, erbB3 and erbB4, are frequently involved in driving the proliferation and survival of tumour cells (reviewed in Olayioye *et al.*, EMBO J., 2000, 19, 3159). One mechanism in which this can be accomplished is by overexpression of the receptor at the protein level, generally as a result of gene amplification. This has been observed in many common human cancers (reviewed in Klapper *et al.*, Adv. Cancer Res., 2000, 77, 25) such as breast cancer (Sainsbury *et al.*, Brit. J. Cancer, 1988, 58, 458; Guerin *et al.*, Oncogene Res., 1988, 3, 21; Slamon *et al.*, Science, 1989, 244, 707; Klijn *et al.*, Breast Cancer Res. Treat., 1994, 29, 73 and reviewed in Salomon *et al.*, Crit. Rev. Oncol. Hematol., 1995, 19, 183), non-small cell lung cancers (NSCLCs) including adenocarcinomas (Cerny *et al.*, Brit. J. Cancer, 1986, 54,

265; Reubi *et al.*, *Int. J. Cancer*, 1990, 45, 269; Rusch *et al.*, *Cancer Research*, 1993, 53, 2379; Brabender *et al.*, *Clin. Cancer Res.*, 2001, 7, 1850) as well as other cancers of the lung (Hendler *et al.*, *Cancer Cells*, 1989, 7, 347; Ohsaki *et al.*, *Oncol. Rep.*, 2000, 7, 603), bladder cancer (Neal *et al.*, *Lancet*, 1985, 366; Chow *et al.*, *Clin. Cancer Res.*, 2001, 7, 1957, Zhau *et al.*, *Mol Carcinog.*, 3, 254), oesophageal cancer (Mukaida *et al.*, *Cancer*, 1991, 68, 142), gastrointestinal cancer such as colon, rectal or stomach cancer (Bolen *et al.*, *Oncogene Res.*, 1987, 1, 149; Kapitanovic *et al.*, *Gastroenterology*, 2000, 112, 1103; Ross *et al.*, *Cancer Invest.*, 2001, 19, 554), cancer of the prostate (Visakorpi *et al.*, *Histochem. J.*, 1992, 24, 481; Kumar *et al.*, 2000, 32, 73; Scher *et al.*, *J. Natl. Cancer Inst.*, 2000, 92, 1866), leukaemia  
10 (Konaka *et al.*, *Cell*, 1984, 37, 1035, Martin-Subero *et al.*, *Cancer Genet Cytogenet.*, 2001, 127, 174), ovarian (Hellstrom *et al.*, *Cancer Res.*, 2001, 61, 2420), head and neck (Shiga *et al.*, *Head Neck*, 2000, 22, 599) or pancreatic cancer (Ovotny *et al.*, *Neoplasma*, 2001, 48, 188). As more human tumour tissues are tested for expression of the erbB family of receptor tyrosine kinases it is expected that their widespread prevalence and importance will be further  
15 enhanced in the future.

As a consequence of the mis-regulation of one or more of these receptors, it is widely believed that many tumours become clinically more aggressive and so correlate with a poorer prognosis for the patient (Brabender *et al.*, *Clin. Cancer Res.*, 2001, 7, 1850; Ross *et al.*, *Cancer Investigation*, 2001, 19, 554, Yu *et al.*, *Bioessays*, 2000, 22.7, 673). In addition to these  
20 clinical findings, a wealth of pre-clinical information suggests that the erbB family of receptor tyrosine kinases are involved in cellular transformation. This includes the observations that many tumour cell lines overexpress one or more of the erbB receptors and that EGFR or erbB2 when transfected into non-tumour cells have the ability to transform these cells. This tumourigenic potential has been further verified as transgenic mice that overexpress erbB2  
25 spontaneously develop tumours in the mammary gland. In addition to this, a number of pre-clinical studies have demonstrated that anti-proliferative effects can be induced by knocking out one or more erbB activities by small molecule inhibitors, dominant negatives or inhibitory antibodies (reviewed in Mendelsohn *et al.*, *Oncogene*, 2000, 19, 6550). Thus it has been recognised that inhibitors of these receptor tyrosine kinases should be of value as a  
30 selective inhibitor of the proliferation of mammalian cancer cells (Yaish *et al.* *Science*, 1988, 242, 933, Kolibaba *et al.*, *Biochimica et Biophysica Acta*, 1997, 133, F217-F248; Al-Obeidi *et al.*, 2000, *Oncogene*, 19, 5690-5701; Mendelsohn *et al.*, 2000, *Oncogene*, 19, 6550-6565). In addition to this pre-clinical data, findings using inhibitory antibodies against EGFR and erbB2

- 4 -

(c-225 and trastuzumab respectively) have proven to be beneficial in the clinic for the treatment of selected solid tumours (reviewed in Mendelsohn *et al.*, 2000, Oncogene, 19, 6550-6565).

Amplification and/or activity of members of the erbB type receptor tyrosine kinases have been detected and so have been implicated to play a role in a number of non-malignant proliferative disorders such as psoriasis (Ben-Bassat, Curr. Pharm. Des., 2000, 6, 933; Elder *et al.*, Science, 1989, 243, 811), benign prostatic hyperplasia (BPH) (Kumar *et al.*, Int. Urol. Nephrol., 2000, 32, 73), atherosclerosis and restenosis (Bokemeyer *et al.*, Kidney Int., 2000, 58, 549). It is therefore expected that inhibitors of erbB type receptor tyrosine kinases will be useful in the treatment of these and other non-malignant disorders of excessive cellular proliferation.

European patent application EP 566 226 discloses certain 4-anilinoquinazolines that are receptor tyrosine kinase inhibitors.

International patent applications WO 96/33977, WO 96/33978, WO 96/33979, WO 96/33980, WO 96/33981, WO 97/30034 and WO 97/38994 disclose that certain quinazoline derivatives which bear an anilino substituent at the 4-position and a substituent at the 6- and/or 7- position possess receptor tyrosine kinase inhibitory activity.

European patent application EP 837 063 discloses aryl substituted 4-aminoquinazoline derivatives carrying a moiety containing an aryl or heteroaryl group at the 6- or 7- position on the quinazoline ring. The compounds are stated to be useful for treating hyperproliferative disorders.

International patent applications WO 97/30035 and WO 98/13354 disclose certain 4-anilinoquinazolines substituted at the 7- position are vascular endothelial growth factor receptor tyrosine kinase inhibitors.

WO 00/55141 discloses 6,7-substituted 4-anilinoquinazoline compounds characterised in that the substituents at the 6- and/or 7- position carry an ester linked moiety (RO-CO).

WO 00/56720 discloses 6,7-dialkoxy-4-anilinoquinazoline compounds for the treatment of cancer or allergic reactions.

WO 02/41882 discloses 4-anilinoquinazoline compounds substituted at the 6- and/or 7- position by a substituted pyrrolidinyloxy or piperidinyloxy group.

WO 03/082290 discloses that certain 6,7-substituted 4-anilinoquinazoline compounds possess receptor tyrosine kinase inhibitory activity. A specific example of such a compound is 6-[[1-(N-methylcarbamoylmethyl)piperidin-4-yl]oxy]-4-(3-chloro-4-fluoroanilino)

- 5 -

-7-methoxy-quinazoline.


None of the prior art discloses 4-(2,3-dihalogenoanilino)quinazoline or 4-(2,3,4-trihalogenoanilino)quinazoline compounds.

Copending International Patent Application No. PCT/GB03/01306 describes that certain 4-(2,3-dihalogenoanilino)quinazoline derivatives possess potent anti-tumour activity, and in particular are selective against EGFR. A specific example of such a compound is 6-[[1-(carbamoylmethyl)piperidin-4-yl]methoxy]-4-(3-chloro-2-fluoroanilino)-7-methoxy-quinazoline.

The applicants have surprisingly found however that addition of a substituent to the carbamoyl group and, optionally, adding a further substituent to the aniline group produces a select group of compounds with enhanced activity in that the compounds have a dual activity, being particularly effected as erbB2 kinase inhibitors, whilst retaining the EGF inhibitory effect, making them of particular clinical application in the treatment of tumours where both these kinases are implicated.

Without wishing to imply that the compounds disclosed in the present invention possess pharmacological activity only by virtue of an effect on a single biological process, it is believed that the compounds provide an anti-tumour effect by way of inhibition of two of the erbB family of receptor tyrosine kinases that are involved in the signal transduction steps which lead to the proliferation of tumour cells. In particular, it is believed that the compounds of the present invention provide an anti-tumour effect by way of inhibition of EGFR and/or erbB2 receptor tyrosine kinases.

According to a first aspect of the invention there is provided a quinazoline derivative of the Formula I:



wherein *n* is 0, 1, 2 or 3,

- 6 -

- each  $R^5$  is independently selected from halogeno, cyano, nitro, hydroxy, amino, carboxy, sulfamoyl, trifluoromethyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, 5 N-(1-6C)alkylsulfamoyl, and N,N-di-[(1-6C)alkyl]sulfamoyl,  $C(O)NR^6R^7$  where  $R^6$  and  $R^7$  are independently selected from hydrogen, optionally substituted (1-6C)alkyl, optionally substituted (3-8C)cycloalkyl or optionally substituted aryl, or  $R^6$  and  $R^7$  together with the nitrogen to which they are attached form an optionally substituted heterocyclic ring which may contain additional heteroatoms;
- 10  $X^1$  is a direct bond or O;  
 $R^1$  is selected from hydrogen and (1-6C)alkyl, wherein the (1-6C)alkyl group is optionally substituted by one or more substituents, which may be the same or different, selected from hydroxy and halogeno, and/or a substituent selected from amino, nitro, carboxy, cyano, halogeno, (1-6C)alkoxy, hydroxy(1-6C)alkoxy, (2-8C)alkenyl, (2-8C)alkynyl, 15 (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, carbamoyl, N-(1-6C)alkylcarbamoyl, N,N di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, (1-6C)alkoxycarbonyl, sulfamoyl, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and 20 N-(1-6C)alkyl-(1-6C)alkanesulfonylamino;  
 $m$  is 0, 1, 2 or 3;  
 $R^2$  is hydrogen or (1-6C)alkyl; and  
 $R^3$  is (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl or (1-6C)alkoxy, any of which can be optionally substituted on a carbon atom by a (1-6C)alkoxy, amino, (1-6C)alkylamino, 25 di-(1-6C)alkylamino, or a group  $S(O)_s(1-6C)alkyl$  where  $s$  is 0, 1 or 2, or a saturated 5 or 6 membered heterocyclic ring which optionally contains additional heteroatoms selected from oxygen, sulfur or  $NR^8$  where  $R^8$  is hydrogen, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkylsulfonyl or (1-6C)alkylcarbonyl;  
or  $R^2$  and  $R^3$  together with the nitrogen atom to which they are attached form a saturated 5 or 30 6 membered heterocyclic ring which optionally contains additional heteroatoms selected from oxygen, S, SO or  $S(O)_2$  or  $NR^8$  where  $R^8$  is as defined above;  
provided that the quinazoline derivative is not:



- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 5 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy}-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonyl]-piperidin-4-yl-oxy}-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(diethylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 10 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(piperidin-1-yl)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(pyrrolidin-1-yl)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 15 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(4-methyl-piperazin-1-yl)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy}-7-ethoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy}-7-(2-methoxy-ethoxy)-quinazoline;
- 20 4-[(3-ethynyl-phenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(ethylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 25 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(isopropylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonylmethyl]-piperidin-4-yl-oxy}-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonylmethyl]-piperidin-4-yl-oxy}-quinazoline;
- 30 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;

- 8 -

- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(methylamino)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 5 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(pyrrolidin-1-yl)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 10 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(methylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(2-methoxyethyl)amino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 15 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(N-methyl-N-2-methoxyethyl)amino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(3-methoxypropyl)amino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(N-methyl-N-3-methoxypropyl)amino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 20 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonylethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline; or
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonylpropyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 25 or a pharmaceutically acceptable salt thereof.

In this specification the generic term "alkyl" includes both straight-chain and branched-chain alkyl groups such as propyl, isopropyl and tert-butyl, and (3-7C)cycloalkyl groups such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. However references to individual alkyl groups such as "propyl" are specific for the straight-chain

30 version only, references to individual branched-chain alkyl groups such as "isopropyl" are specific for the branched-chain version only and references to individual cycloalkyl groups such as "cyclopentyl" are specific for that 5-membered ring only. An analogous convention applies to other generic terms, for example (1-6C)alkoxy includes methoxy, ethoxy,

- 9 -

cyclopropyloxy and cyclopentyloxy, (1-6C)alkylamino includes methylamino, ethylamino, cyclobutylamino and cyclohexylamino, and di-[(1-6Calkyl)]amino includes dimethylamino, diethylamino, N-cyclobutyl-N-methylamino and N-cyclohexyl-N-ethylamino.

The term "aryl" refers to aromatic hydrocarbon rings such as phenyl or naphthyl. The terms "heterocyclic" or "heterocyclyl" include ring structures that may be mono- or bicyclic and contain from 3 to 15 atoms, at least one of which, and suitably from 1 to 4 of which, is a heteroatom such as oxygen, sulfur or nitrogen. Rings may be aromatic, non-aromatic or partially aromatic in the sense that one ring of a fused ring system may be aromatic and the other non-aromatic. Particular examples of such ring systems include furyl, benzofuranyl, tetrahydrofuryl, chromanyl, thienyl, benzothienyl, pyridyl, piperidyl, quinolyl, 1,2,3,4-tetrahydroquinolyl, isoquinolyl, 1,2,3,4-tetrahydroisoquinolyl, pyrazinyl, piperazinyl, pyrimidinyl, pyridazinyl, quinoxalyl, quinazolyl, cinnolyl, pyrrolyl, pyrrolidinyl, indolyl, indolyl, imidazolyl, benzimidazolyl, pyrazolyl, indazolyl, oxazolyl, benzoxazolyl, isoxazolyl, thiazolyl, benzothiazolyl, isothiazolyl, morpholyl, 4H-1,4-benzoxazinyl, 4H-1,4-benzothiazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, oxadiazolyl, furazanyl, thiadiazolyl, tetrazolyl, dibenzofuranyl, dibenzothienyl oxiranyl, oxetanyl, azetidyl, tetrahydropyranyl, oxepanyl, oxazepanyl, tetrahydro-1,4-thiazinyl, 1,1-dioxotetrahydro-1,4-thiazinyl, homopiperidinyl, homopiperazinyl, dihydropyridinyl, tetrahydropyridinyl, dihydropyrimidinyl, tetrahydropyrimidinyl, tetrahydrothienyl, tetrahydrothiopyranyl or thiomorpholyl.

Particular examples of heterocyclic groups include tetrahydropyranyl, tetrahydrofuryl or N-(1-6C)alkylpyrrolidine or N-alkyl(1-6C)piperidine.

Where rings include nitrogen atoms, these may carry a hydrogen atom or a substituent group such as an (1-6C)alkyl group if required to fulfil the bonding requirements of nitrogen, or they may be linked to the rest of the structure by way of the nitrogen atom. A nitrogen atom within a heterocyclyl group may be oxidized to give the corresponding N oxide.

Generally the compounds exhibit favourable physical properties such as a high solubility whilst retaining high antiproliferative activity. Furthermore, many of the compounds according to the present invention are inactive or only weakly active in a hERG assay.

It is to be understood that, insofar as certain of the compounds of Formula I defined above may exist in optically active or racemic forms by virtue of one or more asymmetrically substituted carbon and/or sulfur atoms, and accordingly may exist in, and be isolated as

- 10 -

enantiomerically pure, a mixture of diastereoisomers or as a racemate. The present invention includes in its definition any racemic, optically-active, enantiomerically pure, mixture of diastereoisomers, stereoisomeric form of the compound of Formula I, or mixtures thereof, which possesses the above-mentioned activity. The synthesis of optically active forms may  
5 be carried out by standard techniques of organic chemistry well known in the art, for example by synthesis from optically active starting materials or by resolution of a racemic form. Similarly, the above-mentioned activity may be evaluated using the standard laboratory techniques referred to hereinafter.

The invention relates to all tautomeric forms of the compounds of the Formula I that  
10 possess antiproliferative activity.

It is also to be understood that certain compounds of the Formula I may exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which possess antiproliferative activity.

15 It is also to be understood that certain compounds of the Formula I may exhibit polymorphism, and that the invention encompasses all such forms which possess antiproliferative activity.

Suitable values for the generic radicals referred to above include those set out below.

Suitable values for any of the R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> or R<sup>5</sup> groups as defined hereinbefore or  
20 hereafter in this specification include:-

for halogeno	fluoro, chloro, bromo and iodo;
for (1-6C)alkyl:	methyl, ethyl, propyl, isopropyl, <u>tert</u> -butyl, pentyl and hexyl;
for (1-4C)alkyl:	methyl, ethyl, propyl, isopropyl and <u>tert</u> -butyl;
25 for (1-6C)alkoxy:	methoxy, ethoxy, propoxy, isopropoxy and butoxy;
for (2-8C)alkenyl:	vinyl, isopropenyl, allyl and but-2-enyl;
for (2-8C)alkynyl:	ethynyl, 2-propynyl and but-2-ynyl;
for (2-6C)alkenyloxy:	vinylloxy and allyloxy;
for (2-6C)alkynyloxy:	ethynyloxy and 2-propynyloxy;
30 for (1-6C)alkylthio:	methylthio, ethylthio and propylthio;
for (1-6C)alkylsulfinyl:	methylsulfinyl and ethylsulfinyl;
for (1-6C)alkylsulfonyl:	methylsulfonyl and ethylsulfonyl;

- 11 -

- for (1-6C)alkylamino: methylamino, ethylamino, propylamino,  
isopropylamino and butylamino;
- for di-[(1-6C)alkyl]amino: dimethylamino, diethylamino, N-ethyl-  
N-methylamino and diisopropylamino;
- 5 for (1-6C)alkoxycarbonyl: methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl  
and tert-butoxycarbonyl;
- for N-(1-6C)alkylcarbamoyl: N-methylcarbamoyl, N-ethylcarbamoyl,  
N-propylcarbamoyl and N-isopropylcarbamoyl;
- for N,N-di-[(1-6C)alkyl]carbamoyl: N,N-dimethylcarbamoyl, N-ethyl-  
10 N-methylcarbamoyl and N,N-diethylcarbamoyl;
- for (2-6C)alkanoyl: acetyl, propionyl and isobutyryl;
- for (2-6C)alkanoyloxy: acetoxyl and propionyloxy;
- for (2-6C)alkanoylamino: acetamido and propionamido;
- for N-(1-6C)alkyl-(2-6C)alkanoylamino: N-methylacetamido and N-methylpropionamido;
- 15 for N-(1-6C)alkylsulfamoyl: N-methylsulfamoyl, N-ethylsulfamoyl and  
N-isopropylsulfamoyl;
- for N,N-di-[(1-6C)alkyl]sulfamoyl: N,N-dimethylsulfamoyl and  
N-methyl-N-ethylsulfamoyl;
- for (1-6C)alkanesulfonylamino: methanesulfonylamino and ethanesulfonylamino;
- 20 for N-(1-6C)alkyl-(1-6C)alkanesulfonylamino: N-methylmethanesulfonylamino and  
N-methylethanesulfonylamino;
- for hydroxy-(1-6C)alkoxy: hydroxymethoxy, 2-hydroxyethoxy,  
1-hydroxyethoxy and 3-hydroxypropoxy.

It is to be understood that when R<sup>1</sup> is a group (1-6C)alkyl substituted by, for example  
25 amino to give for example a 2-aminoethyl group, it is the (1-6C)alkyl group that is attached to  
the group X<sup>1</sup> (or the quinazoline ring when X<sup>1</sup> is a direct bond).

When in this specification reference is made to a (1-4C)alkyl group it is to be  
understood that such groups refer to alkyl groups containing up to 4 carbon atoms. A skilled  
person will realise that representative examples of such groups are those listed above under  
30 (1-6C)alkyl that contain up to 4 carbon atoms, such as methyl, ethyl, propyl, isopropyl, butyl  
and tert-butyl. Similarly, reference to a (1-3C)alkyl group refers to alkyl groups containing  
up to 3 carbon atoms such as methyl, ethyl, propyl and isopropyl. A similar convention is

- 12 -

adopted for the other groups listed above such as (1-4C)alkoxy, (2-4C)alkenyl, (2-4C)alkynyl and (2-4C)alkanoyl.

In the compound of Formula I hydrogen atoms are present at the 2, 5 and 8 positions on the quinazoline ring.

5 A suitable pharmaceutically-acceptable salt of a compound of the Formula I is, for example, an acid-addition salt of a compound of the Formula I, for example an acid-addition salt with an inorganic or organic acid such as hydrochloric, hydrobromic, sulfuric, trifluoroacetic, citric or maleic acid; or, for example, a salt of a compound of the Formula I which is sufficiently acidic, for example an alkali or alkaline earth metal salt such as a  
10 calcium or magnesium salt, or an ammonium salt, or a salt with an organic base such as methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

Particular examples of  $n$  are 1, 2 or 3, suitably 2 or 3.

Suitably each  $R^5$  is independently selected from halogeno, trifluoromethyl,  
15 (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl or a group  $C(O)NR^6R^7$  where  $R^6$  and  $R^7$  are as defined above.

In particular, each group  $R^5$  is independently selected from halogeno, such as chloro or fluoro.

Particular substituents for groups  $R^6$  and  $R^7$  where these are other than hydrogen,  
20 include halogeno, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, sulfamoyl, trifluoromethyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfanyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkyl carbamoyl, N,N-di-[(1-6C)alkyl] carbamoyl, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl,  
25 (3-8C)cycloalkyl, aryl or heterocyclic groups.

Particular examples of aryl substituents for  $R^6$  or  $R^7$  include phenyl or naphthyl, particularly phenyl.

Particular examples of heterocyclic substituents for  $R^6$  or  $R^7$  include 5 or 6 membered heterocyclic rings such as furyl, tetrahydrofuryl, thienyl, pyridyl, piperidinyl, pyrazinyl,  
30 piperazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, pyrrolidinyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, morpholinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, oxadiazolyl, furazanyl, thiadiazolyl or tetrazolyl.

- 13 -

When R<sup>6</sup> and R<sup>7</sup> together with the nitrogen to which they are attached form an optionally substituted heterocyclic ring, it is for example a 5 or 6 membered ring, which is saturated or unsaturated. Particular examples include piperidinyl, pyrrolidinyl, morpholinyl or thiomorpholino. Alternatively, R<sup>6</sup> and R<sup>7</sup> together form a (3-6C)alkenyl group.

5 Heterocyclic rings formed by R<sup>6</sup> and R<sup>7</sup> together with the nitrogen atom to which they are attached may be substituted by any of the groups mentioned above in relation to R<sup>6</sup> and R<sup>7</sup>. In addition, these rings may be substituted by one or more (1-6C) alkyl groups, which may themselves be optionally substituted by one or more groups selected from halogeno, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, sulfamoyl, trifluoromethyl, (2-8C)alkenyl, 10 (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, N-(1-6C)alkylsulfamoyl, or N,N-di-[(1-6C)alkyl]sulfamoyl.

An exemplary group of substituents for R<sup>6</sup> or R<sup>7</sup> where they are other than hydrogen 15 are cyano, hydroxy, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylamino, aryl such as phenyl or heterocyclic groups such as furyl, and additionally, where R<sup>6</sup> and R<sup>7</sup> together with the nitrogen atom to which they are attached form a ring, (1-6C) alkyl groups such as methyl.

Where n is 1, 2 or 3, one group R<sup>5</sup> is suitably at an ortho- (2-) position on the benzene 20 ring.

Where n is 1, 2 or 3, one group R<sup>5</sup> is suitably at a meta- (3-) position on the benzene ring.

Thus, when n is 1, the R<sup>5</sup> group is suitably at an ortho- (2-) or a meta- (3-) position on the benzene ring.

25 In one aspect of the invention, when n is 2, the first R<sup>5</sup> group is suitably at a meta-position and the second R<sup>5</sup> group is suitably at an ortho- or a para- position on the benzene ring, and thus the ring has substituents at 2- and 3- or 3- and 4- positions on the benzene ring.

In another aspect of the invention, when n is 2 or 3, the first R<sup>5</sup> group is suitably at an ortho- position, the second R<sup>5</sup> group is suitably at a meta- position and, optionally (when n is 30 3), the third R<sup>5</sup> group is suitably at a para- position on the benzene ring. Thus, when n is 2, the ring suitably has substituents at 2- and 3- positions on the benzene ring and when n is 3, the ring suitably has substituents at 2-, 3- and 4- positions on the benzene ring.

- 14 -


The applicants have surprisingly found that quinazoline derivatives having substituents (for example halogeno substituents) at 2- and 3- positions or at 2-, 3- and 4- positions on the benzene ring compared to quinazoline derivatives having substituents at 3- and 4- positions on the benzene ring produces compounds with enhanced activity in that the compounds have an increased potency against erbB2 and/or EGFR (particularly erbB2) receptor tyrosine kinases in cellular assays. It is believed that quinazoline derivatives having substituents (for example halogeno substituents) at 2- and 3- positions or at 2-, 3- and 4- positions on the benzene ring will also have an increased potency against erbB2 and/or EGFR (particularly erbB2) receptor tyrosine kinases *in vivo*.

10 Suitably when n is 2 or 3, each R<sup>5</sup> group is the same or different halogeno atom, such as chloro or fluoro. Suitably, at least one R<sup>5</sup> group is fluoro, which at least one fluoro is preferably positioned at an ortho- (2-) position on the benzene ring.


Suitably when n is 2, each R<sup>5</sup> group is the same or different halogeno atom. In particular, one R<sup>5</sup> group is chloro, and this is preferably at a meta- (3-) position on the benzene ring to which it is attached, and the other R<sup>5</sup> group is fluoro which is preferably at an ortho- (2-) or a para- (4-) (preferably an ortho- (2-)) position on the benzene ring.

Suitably when n is 3, each R<sup>5</sup> group is the same or different halogeno atom. In particular, one R<sup>5</sup> group is chloro, and this is preferably at a meta- (3-) position on the benzene ring to which it is attached, and the other two R<sup>5</sup> groups are each fluoro, which are preferably at an ortho- (2-) and a para- (4-) position respectively on the benzene ring.

Thus particular examples of the group of sub-formula (i):



in Formula I are groups of sub-formula (ii):



25 wherein (a) one of R<sup>10</sup> or R<sup>12</sup> is hydrogen and the other is halogeno, such as chloro or fluoro, and particularly fluoro, and R<sup>11</sup> is halogeno such as chloro or fluoro, and particularly



- 15 -

chloro, or (b)  $R^{10}$  is halogeno, such as chloro or fluoro, and particularly fluoro,  $R^{11}$  is halogeno such as chloro or fluoro, and particularly chloro, and  $R^{12}$  is hydrogen or halogeno, such as chloro or fluoro, and particularly fluoro, or (c)  $R^{10}$  is fluoro,  $R^{11}$  is chloro, and  $R^{12}$  is hydrogen or fluoro. In particular,  $R^{10}$ ,  $R^{11}$  and  $R^{12}$  are as defined in (b) and/or (c).

- 5 In one embodiment, when  $n$  is 2, each  $R^5$  group is the same or different halogeno atom (such as fluoro and/or chloro) and the first  $R^5$  group is at an ortho- position and the second  $R^5$  group is at a meta- position on the benzene ring, then (i) when  $m$  is 0, 1, 2 or 3,  $R^3$  is not (1-6C)alkyl and (ii) when  $m$  is 0,  $R^2$  and  $R^3$  do not, together with the nitrogen atom to which they are attached, form a saturated 5 or 6 membered heterocyclic ring which optionally contains  
10 additional heteroatoms selected from oxygen, S, SO or  $S(O)_2$  or  $NR^8$  where  $R^8$  is hydrogen, (1-4C)alkyl or (1-4C)alkylsulfonyl.

Suitably  $X^1$  is oxygen.

- In particular  $R^1$  is selected from hydrogen, (1-6C)alkyl and (1-6C)alkoxy(1-6C)alkyl, wherein any (1-6C)alkyl group in  $R^1$  optionally bears one or more (suitably 1 or 2) hydroxy or  
15 halogeno substituents. More particularly,  $R^1$  is selected from (1-6C)alkyl, preferably from (1-4C)alkyl and even more preferably from (1-2C)alkyl. For example,  $R^1$  may be methyl.

For instance,  $R^1-X^1$  is selected from methoxy, ethoxy, isopropoxy, cyclopropylmethoxy, 2-hydroxyethoxy, 2-fluoroethoxy, 2-methoxyethoxy, 2,2-difluoroethoxy, 2,2,2-trifluoroethoxy or 3-hydroxy-3-methylbutoxy.

- 20 In particular  $R^1-X$  is selected from hydrogen, (1-4C)alkoxy and (1-4C)alkoxy(1-4C)alkoxy. For instance,  $R^1-X$  is selected from hydrogen, methoxy, ethoxy and 2-methoxyethoxy. A particular example of a group  $R^1-X^1$  is methoxy.

Suitably  $m$  is 1, 2 or 3. Preferably  $m$  is 0 or 1 (more preferably 1).

- When  $R^2$  and  $R^3$  together with the nitrogen atom to which they are attached form a  
25 saturated heterocyclic ring which optionally contains additional heteroatoms, the heterocyclic ring is particularly a 6 membered ring.

- When  $R^2$  and  $R^3$  together with the nitrogen atom to which they are attached form a saturated 5 or 6 (preferably 6) membered heterocyclic ring which optionally contains additional heteroatoms, this suitably contains additional heteroatoms selected from O and  
30  $NR^8$ , where  $R^8$  is as defined in relation to Formula I.

When  $R^2$  and  $R^3$  together with the nitrogen atom to which they are attached form a saturated 5 or 6 membered heterocyclic ring which optionally contains additional heteroatoms, this suitably comprises a pyrrolidine ring, a morpholine ring, a piperidine ring,

- 16 -

or a piperazine ring which is optionally substituted on the available nitrogen atom by a group  $R^8$  as defined above. In particular, the heterocyclic ring comprises a morpholine ring or a piperazine ring which is optionally substituted on the available nitrogen atom by a group  $R^8$  as defined in relation to Formula I.

- 5 Particular examples of  $R^8$  groups include (1-3C) alkyl such as methyl; (1-3C)alkylsulfonyl such as methylsulfonyl; (1-3C)alkylcarbonyl, such as acetyl; (2-4C)alkenyl such as allyl; or (2-4C)alkynyl such as propargyl. In particular  $R^8$  is a (1-3C)alkyl group such as methyl.

- Thus when  $R^2$  together with  $R^3$  together with the nitrogen atom to which they are  
10 attached form a saturated 5 or 6 membered heterocyclic ring which optionally contains additional heteroatoms, this suitably comprises a morpholine ring. Other examples include pyrrolidine, piperidine, piperazine or N-methylpiperazine, particularly piperazine or N-methylpiperazine.

Preferably  $R^2$  is hydrogen or (1-3C)alkyl.


- 15 In particular  $R^2$  is hydrogen or methyl (preferably hydrogen).

Suitable substituents for  $R^3$  include (1-6C)alkoxy, (1-6C)alkylamino, di-(1-6C)alkylamino or a saturated 5 or 6 membered heterocyclic ring which optionally contains additional heteroatoms selected from oxygen, sulfur or  $NR^8$  where  $R^8$  is as defined above.

- 20 In particular, suitable substituents for  $R^3$  include (1-3C)alkoxy such as methoxy, amino, (1-3C)alkylamino, di-(1-3C)alkylamino such as dimethylamino, (1-3C)alkylsulfonyl, a pyrrolidine ring or a piperazine ring, which may contain on the available nitrogen atom a (1-3C)alkyl group such as methyl, a (2-4C)alkenyl group such as vinyl, an (2-4C)alkynyl group such as propargyl, an (1-5C)alkylsulfonyl group such as methyl sulfonyl or an  
25 (1-6C)alkylcarbonyl group such as acetyl.

Suitably  $R^3$  is (1-6C)alkyl, in particular (1-3C)alkyl, such as methyl or ethyl. Suitably  $R^2$  is hydrogen and  $R^3$  is (1-6C)alkyl, in particular (1-3C)alkyl, such as methyl or ethyl.

Particular examples of the compounds of Formula I are compounds of Formula IA:



IA

where  $R^2$ ,  $R^3$  and  $m$  are as defined above,  $R^{10}$ ,  $R^{11}$  and  $R^{12}$  are as defined in relation to sub-formula (ii) above, and  $R^{13}$  is selected from hydrogen, methoxy, ethoxy and 2-methoxyethoxy, and especially methoxy.

5 For the avoidance of any doubt, when the compounds of Formula I are defined as compounds of Formula IA, the quinazoline derivative is not:

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy}-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonyl]-piperidin-4-yl-oxy}-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(diethylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(piperidin-1-yl)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(pyrrolidin-1-yl)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(4-methyl-piperazin-1-yl)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy}-7-ethoxy-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy}-7-(2-methoxy-ethoxy)-quinazoline;


- 18 -

- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(ethylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(isopropylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 5 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonylmethyl]-piperidin-4-yl-oxy}-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonylmethyl]-piperidin-4-yl-oxy}-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 10 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(methylamino)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 15 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(pyrrolidin-1-yl)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 20 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(methylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(2-methoxyethyl)amino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 25 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(N-methyl-N-2-methoxyethyl)amino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(3-methoxypropyl)amino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(N-methyl-N-3-methoxypropyl)amino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 30 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]ethyl}-piperidin-4-yl-oxy}-7-methoxy-quinazoline; or


- 19 -

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonylpropyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;  
or a pharmaceutically acceptable salt thereof.

Other particular examples of the compounds of Formula I are compounds of the  
5 Formulae IB and IC:




IB



IC

where  $R^2$ ,  $R^3$  and  $m$  are as defined above and  $R^{13}$  is selected from hydrogen, methoxy, ethoxy and 2-methoxyethoxy, and especially methoxy.

10 Other particular examples of the compounds of Formula I are compounds of the  
Formula ID:



ID

wherein:

- 20 -

$R^{5a}$  and  $R^{5b}$  are independently selected from halogeno (for example fluoro and/or chloro);

$X^1$  is a direct bond or O;

$R^1$  is selected from hydrogen and (1-6C)alkyl, wherein the (1-6C)alkyl group is optionally substituted by one or more substituents, which may be the same or different, selected from

5 hydroxy and halogeno, and/or a substituent selected from amino, nitro, carboxy, cyano, halogeno, (1-6C)alkoxy, hydroxy(1-6C)alkoxy, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, carbamoyl, N-(1-6C)alkylcarbamoyl, N,N di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino,

10 N-(1-6C)alkyl-(2-6C)alkanoylamino, (1-6C)alkoxycarbonyl, sulfamoyl,

N-(1-6C)alkylsulfamoyl, N, N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and

N-(1-6C)alkyl-(1-6C)alkanesulfonylamino;

$m$  is 0, 1, 2 or 3;

$R^2$  is hydrogen or (1-6C)alkyl; and

15  $R^3$  is (1-6C)alkyl, wherein the (1-6C)alkyl group is optionally substituted on a carbon atom by a (1-6C)alkoxy, amino, (1-6C)alkylamino, di-(1-6C)alkylamino, or a group  $S(O)_s(1-6C)alkyl$  where  $s$  is 0, 1 or 2, or a saturated 5 or 6 membered heterocyclic ring which optionally contains additional heteroatoms selected from oxygen, sulfur or  $NR^8$  where  $R^8$  is hydrogen, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkylsulfonyl or (1-6C)alkylcarbonyl;

20 or when  $m$  is 0,  $R^2$  and  $R^3$  together with the nitrogen atom to which they are attached form a saturated 5 or 6 membered heterocyclic ring which optionally contains additional heteroatoms selected from oxygen, S, SO or  $S(O)_2$  or  $NR^8$  where  $R^8$  is hydrogen, (1-4C)alkyl or (1-4C)alkylsulfonyl;

or a pharmaceutically acceptable salt thereof.

25 In one embodiment of the compounds of the Formula ID, the group  $R^3$  is (1-6C)alkyl, particularly unsubstituted (1-6C)alkyl. For example, the group  $R^3$  may be methyl or ethyl, particularly methyl.

In another embodiment of the compounds of the Formula ID,  $m$  is 0 and  $R^2$  and  $R^3$  together with the nitrogen atom to which they are attached form a saturated 5 or 6 membered  
30 heterocyclic ring which optionally contains additional heteroatoms selected from oxygen, S, SO or  $S(O)_2$  or  $NR^8$  where  $R^8$  is hydrogen, (1-4C)alkyl or (1-4C)alkylsulfonyl. For example,  $R^2$  together with  $R^3$  together with the nitrogen atom to which they are attached may form a

morpholine ring. Other examples include pyrrolidine, piperidine, piperazine or N-methylpiperazine, particularly piperazine or N-methylpiperazine.

It would be clear to a person skilled in the art that the particular novel compounds of the invention include those compounds of the Formula I (including IA, IB, IC and ID) in which, unless otherwise stated, each of  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ ,  $X^1$ ,  $m$  and  $n$  has any of the meanings as hereinbefore defined.

Examples of quinazoline derivatives of the Formula I include one or more of:

- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoylmethyl)piperidin-4-yl}-oxy}quinazoline;
- 10 4-(3-chloro-2-fluoroanilino)-6-{{1-(N,N-dimethylcarbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(morpholin-4-ylcarbonylmethyl)piperidin-4-yl}oxy}-quinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(pyrrolidin-1-ylcarbonyl)piperidin-4-yl}oxy}quinazoline;
- 15 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoyl)piperidin-4-yl}oxy}quinazoline;
- 4-(3-chloro-2-fluoroanilino)-6-{{1-(N-(2-dimethylaminoethyl)carbamoyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;
- 20 4-(3-chloro-2-fluoroanilino)-6-{{1-(N,N-dimethylcarbamoyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(morpholin-4-ylcarbonyl)piperidin-4-yl}oxy}quinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-[2-pyrrolidin-1-ylethyl]carbamoyl)piperidin-4-yl}oxy}quinazoline;
- 25 4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoylmethyl)piperidin-4-yl}oxy}quinazoline;
- 4-(3-chloro-2-fluoroanilino)-6-{{1-(N-ethylcarbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;
- 30 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-[2-(pyrrolidin-1-yl)ethyl]carbamoylmethyl)piperidin-4-yl}oxy}quinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-(2-methoxyethyl)carbamoylmethyl)piperidin-4-yl}oxy}quinazoline;

- 22 -

4-(3-chloro-2-fluoroanilino)-6-{{1-(N-(2-dimethylaminoethyl)carbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-({1-[2-(4-methylpiperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy)quinazoline;

5 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-({1-[2-(piperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy)quinazoline; and

4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-({1-[2-(4-methylpiperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy)quinazoline;

or a pharmaceutically acceptable salt thereof.

10 Particular examples of quinazoline derivatives of the Formula I include one or more of:

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoylmethyl)piperidin-4-yl}oxy}quinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoyl)piperidin-4-yl}oxy}

15 quinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(morpholin-4-ylcarbonyl)piperidin-4-yl}oxy}quinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-[2-pyrrolidin-1-ylethyl]carbamoyl)piperidin-4-yl}oxy}quinazoline;

20 4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoylmethyl)piperidin-4-yl}oxy}quinazoline;

4-(3-chloro-2-fluoroanilino)-6-{{1-(N-ethylcarbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-[2-(pyrrolidin-1-yl)ethyl]carbamoylmethyl)

25 piperidin-4-yl}oxy}quinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-(2-methoxyethyl)carbamoylmethyl)piperidin-4-yl}oxy}quinazoline;

4-(3-chloro-2-fluoroanilino)-6-{{1-(N-(2-dimethylaminoethyl)carbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;

30 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-({1-[2-(4-methylpiperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy)quinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-({1-[2-(piperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy)quinazoline; and



4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-({1-[2-(4-methylpiperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy)quinazoline;  
or a pharmaceutically acceptable salt thereof.

A particular group of examples of quinazoline derivatives of the Formula I includes

5 one or more of:

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoylmethyl)piperidin-4-yl}oxy}quinazoline; and

4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoylmethyl)piperidin-4-yl}oxy}quinazoline;

10 or a pharmaceutically acceptable salt thereof.

A particular group of examples of quinazoline derivatives of the Formula IB includes one or more of:

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoylmethyl)piperidin-4-yl}oxy}quinazoline;

15 4-(3-chloro-2-fluoroanilino)-6-{{1-(N,N-dimethylcarbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(morpholin-4-ylcarbonylmethyl)piperidin-4-yl}oxy}-quinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(pyrrolidin-1-ylcarbonyl)piperidin-4-yl}oxy}

20 quinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoyl)piperidin-4-yl}oxy}quinazoline;

4-(3-chloro-2-fluoroanilino)-6-{{1-(N-(2-dimethylaminoethyl)carbamoyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;

25 4-(3-chloro-2-fluoroanilino)-6-{{1-(N,N-dimethylcarbamoyl)piperidin-4-yl}oxy}7-methoxyquinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(morpholin-4-ylcarbonyl)piperidin-4-yl}oxy}quinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-[2-pyrrolidin-1-ylethyl]carbamoyl)}

30 piperidin-4-yl}oxy}quinazoline;

4-(3-chloro-2-fluoroanilino)-6-{{1-(N-ethylcarbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;

- 24 -

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-[2-(pyrrolidin-1-yl)ethyl]carbamoylmethyl)piperidin-4-yl}oxy}quinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-(2-methoxyethyl)carbamoylmethyl)piperidin-4-yl}oxy}quinazoline;

5 4-(3-chloro-2-fluoroanilino)-6-{{1-(N-(2-dimethylaminoethyl)carbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-[2-(4-methylpiperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy}quinazoline; and

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-[2-(piperazin-1-yl)-2-oxoethyl]piperidin-4-

10 yl}oxy}quinazoline;

or a pharmaceutically acceptable salt thereof.

A particular group of examples of quinazoline derivatives of the Formula IC includes one or more of:

4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoylmethyl)piperidin-4-

15 yl}oxy}quinazoline; and

4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-{{1-[2-(4-methylpiperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy}quinazoline;

or a pharmaceutically acceptable salt thereof.

A particular group of examples of quinazoline derivatives of the Formula ID includes

20 one or more of:

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoylmethyl)piperidin-4-yl}-oxy}quinazoline;

4-(3-chloro-2-fluoroanilino)-6-{{1-(N,N-dimethylcarbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;

25 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(pyrrolidin-1-ylcarbonyl)piperidin-4-yl}oxy}quinazoline;

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoyl)piperidin-4-yl}oxy}quinazoline;

4-(3-chloro-2-fluoroanilino)-6-{{1-(N-(2-dimethylaminoethyl)carbamoyl)piperidin-4-yl}oxy}-

30 7-methoxyquinazoline;

4-(3-chloro-2-fluoroanilino)-6-{{1-(N,N-dimethylcarbamoyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;

- 25 -

- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(morpholin-4-ylcarbonyl)piperidin-4-yl}oxy} quinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-[2-pyrrolidin-1-ylethyl]carbamoyl)piperidin-4-yl}oxy} quinazoline;
- 5 4-(3-chloro-2-fluoroanilino)-6-{{1-(N-ethylcarbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-[2-(pyrrolidin-1-yl)ethyl]carbamoylmethyl)piperidin-4-yl}oxy} quinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-(2-methoxyethyl)carbamoylmethyl)piperidin-4-yl}oxy} quinazoline; and
- 10 4-(3-chloro-2-fluoroanilino)-6-{{1-(N-(2-dimethylaminoethyl)carbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;
- or a pharmaceutically acceptable salt thereof.

- Another particular group of examples of quinazoline derivatives of the Formula ID
- 15 includes one or more of:
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoylmethyl)piperidin-4-yl}oxy} quinazoline;
- 4-(3-chloro-2-fluoroanilino)-6-{{1-(N,N-dimethylcarbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;
- 20 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(pyrrolidin-1-ylcarbonyl)piperidin-4-yl}oxy} quinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoyl)piperidin-4-yl}oxy} quinazoline;
- 4-(3-chloro-2-fluoroanilino)-6-{{1-(N,N-dimethylcarbamoyl)piperidin-4-yl}oxy}-7-methoxy-
- 25 quinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(morpholin-4-ylcarbonyl)piperidin-4-yl}oxy} quinazoline; and
- 4-(3-chloro-2-fluoroanilino)-6-{{1-(N-ethylcarbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;
- 30 or a pharmaceutically acceptable salt thereof.

A preferred example of a compound of the Formula I is

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoylmethyl)piperidin-4-yl}oxy} quinazoline, or a pharmaceutically acceptable salt thereof.

- 26 -

Another preferred example of a compound of the Formula I is

4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoylmethyl)piperidin-4-yl}oxy}quinazoline, or a pharmaceutically acceptable salt thereof.

Another preferred example of a compound of the Formula I is

5 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-[2-(pyrrolidin-1-yl)ethyl]carbamoylmethyl)piperidin-4-yl}oxy}quinazoline, or a pharmaceutically acceptable salt thereof.

Another preferred example of a compound of the Formula I is

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-({1-[2-(4-methylpiperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy)quinazoline, or a pharmaceutically acceptable salt thereof.

10 Another preferred example of a compound of the Formula I is

4-(3-chloro-2-fluoroanilino)-7-methoxy-6-({1-[2-(piperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy)quinazoline, or a pharmaceutically acceptable salt thereof.

#### Synthesis of Quinazoline Derivatives of the Formula I

A further aspect the present invention provides a process for preparing a quinazoline  
15 derivative of Formula I or a pharmaceutically-acceptable salt thereof. It will be appreciated that during certain of the following processes certain substituents may require protection to prevent their undesired reaction. The skilled chemist will appreciate when such protection is required, and how such protecting groups may be put in place, and later removed.

For examples of protecting groups see one of the many general texts on the subject,  
20 for example, 'Protective Groups in Organic Synthesis' by Theodora Green (publisher: John Wiley & Sons). Protecting groups may be removed by any convenient method as described in the literature or known to the skilled chemist as appropriate for the removal of the protecting group in question, such methods being chosen so as to effect removal of the protecting group with minimum disturbance of groups elsewhere in the molecule.

25 Thus, if reactants include, for example, groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group,  
30 for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali

metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulfuric or phosphoric acid or trifluoroacetic acid and an

arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for  
5 example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for  
10 example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium, sodium hydroxide or  
15 ammonia. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed,  
20 for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

Resins may also be used as a protecting group.

The protecting groups may be removed at any convenient stage in the synthesis using  
25 conventional techniques well known in the chemical art.

A quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, may be prepared by any process known to be applicable to the preparation of chemically-related compounds. Such processes, when used to prepare a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, are provided as a  
30 further feature of the invention and are illustrated by the following representative examples. Necessary starting materials may be obtained by standard procedures of organic chemistry (see, for example, Advanced Organic Chemistry (Wiley-Interscience), Jerry March). The preparation of such starting materials is described within the accompanying non-limiting


- 28 -

Examples. Alternatively, necessary starting materials are obtainable by analogous procedures to those illustrated which are within the ordinary skill of an organic chemist. Information on the preparation of necessary starting materials or related compounds (which may be adapted to form necessary starting materials) may also be found in the following Patent and

- 5 Application Publications, the contents of the relevant process sections of which are hereby incorporated herein by reference: WO 94/27965, WO 95/03283, WO 96/33977, WO 96/33978, WO 96/33979, WO 96/33980, WO 96/33981, WO 97/30034, WO 97/38994, WO 01/66099, US 5,252,586, EP 520 722, EP 566 226, EP 602 851 and EP 635 507.


- The present invention also provides that quinazoline derivatives of the Formula I, or  
10 pharmaceutically acceptable salts thereof, can be prepared by a process (a) to (m) as follows (wherein the variables are as defined above unless otherwise stated):

**Process (a)** By reacting a compound of the Formula II:



II

- wherein R¹, X¹, R⁵ and n have any of the meanings defined hereinbefore except that any  
15 functional group is protected if necessary,  
with a compound of the Formula III:



III

- wherein R², R³ and m have any of the meanings defined hereinbefore except that any  
functional group is protected if necessary and Lg is a displaceable group, wherein the reaction  
20 is conveniently performed in the presence of a suitable base,

and whereafter any protecting group that is present is removed by conventional means.

A convenient displaceable group Lg is, for example, a halogeno, alkanesulfonyloxy or arylsulfonyloxy group, for example a chloro, bromo, methanesulfonyloxy,

4-nitrobenzenesulfonyloxy or toluene-4-sulfonyloxy group (suitably a methanesulfonyloxy, 4-nitrobenzenesulfonyloxy or toluene-4-sulfonyloxy group).

The reaction is advantageously carried out in the presence of base. A suitable base is, for example, an organic amine base such as, for example, di-isopropylethylamine, pyridine, 2,6-lutidine, collidine, 4-dimethylaminopyridine, triethylamine, N-methylmorpholine or diazabicyclo[5.4.0]undec-7-ene, or for example, an alkali metal or alkaline earth metal carbonate or hydroxide, for example sodium carbonate, potassium carbonate, cesium carbonate, calcium carbonate, sodium hydroxide or potassium hydroxide. Alternatively such a base is, for example, an alkali metal hydride, for example sodium hydride, an alkali metal or alkaline earth metal amide, for example sodium amide or sodium bis(trimethylsilyl)amide, or a sufficiently basic alkali metal halide, for example cesium fluoride or sodium iodide. The reaction is suitably effected in the presence of an inert solvent or diluent, for example an alkanol or ester such as methanol, ethanol, 2-propanol or ethyl acetate, a halogenated solvent such as methylene chloride, trichloromethane or carbon tetrachloride, an ether such as tetrahydrofuran or 1,4-dioxan, an aromatic hydrocarbon solvent such as toluene, or (suitably) a dipolar aprotic solvent such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidin-2-one or dimethylsulfoxide. The reaction is conveniently effected at a temperature in the range, for example, 10 to 150°C (or the boiling point of the solvent), suitably in the range 20 to 90°C.

A particularly suitable base is cesium fluoride. This reaction is suitably performed in an inert dipolar aprotic solvent such as N,N-dimethylacetamide or N,N-dimethylformamide. The reaction is suitably carried out at a temperature of from 25 to 85°C.


**Process (b)** By modifying a substituent in or introducing a substituent into another quinazoline derivative of Formula I or a pharmaceutically acceptable salt thereof, as hereinbefore defined except that any functional group is protected if necessary, and whereafter any protecting group that is present is removed by conventional means.

Methods for converting substituents into other substituents are known in the art. For example an alkylthio group may be oxidised to an alkylsulfinyl or alkylsulfonyl group, a cyano group reduced to an amino group, a nitro group reduced to an amino group, a hydroxy group alkylated to a methoxy group, a bromo group converted to an alkylthio group, an amino group may be acylated to give an alkanoylamino group (for example by reaction with a suitable acid chloride or acid anhydride) or an alkanoyloxy group may be hydrolysed to a hydroxy group (for example an acetyloxyacetyl group may be converted to a hydroxyacetyl



- 30 -

group). Conveniently, one  $R^1$  group may be converted into another  $R^1$  group as a final step in the preparation of a compound of the Formula I.

**Process (c)** By reacting a compound of Formula IV:



- 5 where  $R^1$ ,  $X^1$ ,  $R^5$  and  $n$  are as defined in relation to Formula I except that any functional group is protected if necessary, with a compound of Formula V or V':



- wherein  $R^2$  and  $R^3$  are as defined above except that any functional group is protected if necessary and  $m'$  is 0, 1, 2 or 3, provided that it is not 0 when  $R^2$  is hydrogen, and  $Lg$  is a  
10 displaceable group (for example halogeno such as chloro or bromo).

- The reactions described above are conveniently performed in the presence of a suitable base (such as those described above in process (a), for example potassium carbonate, triethylamine, 4-dimethylaminopyridine or di-isopropylethylamine) and conveniently in the presence of an inert solvent or diluent (for example the inert solvents and diluents described in  
15 process (a) such as N-methylpyrrolidin-2-one, acetonitrile, N,N-dimethylacetamide, methanol, ethanol or methylene chloride).

- Where  $m'$  is 1, 2 or 3, the reaction is suitably effected in the presence of a source of iodide such as sodium iodide or potassium iodide, and at a temperature in the range, for example, 10 to 150°C (or the boiling point of the solvent), suitably in the range 20 to 90°C.  
20 For example, when  $m$  is 1, the reaction may be conducted using triethylamine as a suitable base, potassium iodide as a suitable iodide source and N,N-dimethylacetamide as a suitable solvent or diluent at a temperature of about 80°C.



- 31 -

Where m is 0 the iodide source is not required and the typical temperature for the reaction is 0°C to room temperature.

Reaction of the compound of Formula IV with a compound of Formula V' is useful for preparing compounds where R<sup>2</sup> is hydrogen and m is 0.

- 5 **Process (d)** By removal of a protecting group from a quinazoline derivative of Formula I, or a pharmaceutically acceptable salt thereof.

Suitable methods for removal of protecting groups are well known and are discussed herein. For example for the production of those compounds of the Formula I wherein R<sup>1</sup> contains a primary or secondary amino group, the cleavage of the corresponding compound of  
10 Formula I wherein R<sup>1</sup> contains a protected primary or secondary amino group.

Suitable protecting groups for an amino group are, for example, any of the protecting groups disclosed hereinbefore for an amino group. Suitable methods for the cleavage of such amino protecting groups are also disclosed hereinbefore. In particular, a suitable protecting group is a lower alkoxycarbonyl group such as a tert-butoxycarbonyl group which may be  
15 cleaved under conventional reaction conditions such as under acid-catalysed hydrolysis, for example in the presence of trifluoroacetic acid.

- Process (e)** By reacting a compound of the Formula II as hereinbefore defined with a compound of the Formula III as defined hereinbefore except Lg is OH under Mitsunobu conditions, and whereafter any protecting group that is present is removed by conventional  
20 means.


Suitable Mitsunobu conditions include, for example, reaction in the presence of a suitable tertiary phosphine and a di-alkylazodicarboxylate in an organic solvent such as THF, or suitably dichloromethane and in the temperature range 0°C to 60°C, but suitably at ambient temperature. A suitable tertiary phosphine includes for example tri-n-butylphosphine or  
25 suitably tri-phenylphosphine. A suitable di-alkylazodicarboxylate includes for example diethyl azodicarboxylate (DEAD) or suitably di-tert-butyl azodicarboxylate. Details of Mitsunobu reactions are contained in Tet. Letts., 31, 699, (1990); The Mitsunobu Reaction, D.L.Hughes, Organic Reactions, 1992, Vol.42, 335-656 and Progress in the Mitsunobu Reaction, D.L.Hughes, Organic Preparations and Procedures International, 1996, Vol.28,  
30 127-164.

**Process (f)** For the preparation of those compounds of the Formula I wherein R<sup>1</sup>-X<sup>1</sup> is a hydroxy group by the cleavage of a quinazoline derivative of the Formula I wherein R<sup>1</sup>-X<sup>1</sup> is a (1-6C)alkoxy group.

- 32 -

The cleavage reaction may conveniently be carried out by any of the many procedures known for such a transformation. The cleavage reaction of a compound of the Formula I wherein  $R^1$  is a (1-6C)alkoxy group may be carried out, for example, by treatment of the quinazoline derivative with an alkali metal (1-6C)alkylsulfide such as sodium ethanethiolate or, for example, by treatment with an alkali metal diarylphosphide such as lithium diphenylphosphide. Alternatively the cleavage reaction may conveniently be carried out, for example, by treatment of the quinazoline derivative with a boron or aluminium trihalide such as boron tribromide, or by reaction with an organic or inorganic acid, for example trifluoroacetic acid. Such reactions are suitably carried out in the presence of a suitable inert solvent or diluent as defined hereinbefore. A preferred cleavage reaction is the treatment of a quinazoline derivative of the Formula I with pyridine hydrochloride. The cleavage reactions are suitably carried out at a temperature in the range, for example, from 10 to 150°C, for example from 25 to 80°C.

**Process (g)** For the preparation of those compounds of the Formula I wherein  $X^1$  is O and  $R^1$  is not hydrogen, by the reaction of a compound of the Formula VI:



VI

wherein  $R^2$ ,  $R^3$ ,  $R^5$ ,  $m$  and  $n$  have any of the meanings defined hereinbefore except that any functional group is protected if necessary, with a compound of the formula  $R^1$ -Lg, wherein  $R^1$  has any of the meanings defined hereinbefore except that it is not hydrogen and except that any functional group is protected if necessary and Lg is a displaceable group, wherein the reaction is conveniently performed in the presence of a suitable base;

and whereafter any protecting group that is present is removed by conventional means.

Suitable displaceable groups, Lg, are as hereinbefore defined for process (a), for example chloro or bromo. The reaction is suitably performed in the presence of a suitable base. Suitable solvents, diluents and bases include, for example those hereinbefore described

in relation to process (a). Alternatively, Lg is a hydroxy group, whereupon the reaction can be effected under Mitsunobu conditions, as described above in relation to process (e).

**Process (h)** For the preparation of those compounds of the Formula I wherein R<sup>1</sup> contains a (1-6C)alkoxy or substituted (1-6C)alkoxy group or a (1-6C)alkylamino or substituted (1-6C)alkylamino group, the alkylation, conveniently in the presence of a suitable base as defined hereinbefore for process (a), of a quinazoline derivative of the Formula I wherein R<sup>1</sup> contains a hydroxy group or a primary or secondary amino group as appropriate.


A suitable alkylating agent is, for example, any agent known in the art for the alkylation of hydroxy to alkoxy or substituted alkoxy, or for the alkylation of amino to alkylamino or substituted alkylamino, for example an alkyl or substituted alkyl halide, for example a (1-6C)alkyl chloride, bromide or iodide or a substituted (1-6C)alkyl chloride, bromide or iodide, conveniently in the presence of a suitable base as defined hereinbefore, in a suitable inert solvent or diluent as defined hereinbefore and at a temperature in the range, for example, 10 to 140°C, conveniently at or near ambient temperature. An analogous procedure may be used to introduce optionally substituted (2-6C)alkanoyloxy, (2-6C)alkanoylamino and (1-6C)alkanesulfonylamino groups into R<sup>1</sup>.

Conveniently for the production of those compounds of the Formula I wherein R<sup>1</sup> contains a (1-6C)alkylamino or substituted (1-6C)alkylamino group, a reductive amination reaction may be employed using formaldehyde or a (2-6C)alkanolaldehyde (for example acetaldehyde or propionaldehyde). For example, for the production of those compounds of the Formula I wherein R<sup>1</sup> contains an N-methyl group, the corresponding compound containing a N-H group may be reacted with formaldehyde in the presence of a suitable reducing agent. A suitable reducing agent is, for example, a hydride reducing agent, for example formic acid, an alkali metal aluminium hydride such as lithium aluminium hydride, or, suitably, an alkali metal borohydride such as sodium borohydride, sodium cyanoborohydride, sodium triethylborohydride, sodium trimethoxyborohydride and sodium triacetoxymethylborohydride. The reaction is conveniently performed in a suitable inert solvent or diluent, for example tetrahydrofuran and diethyl ether for the more powerful reducing agents such as lithium aluminium hydride, and, for example, methylene chloride or a protic solvent such as methanol and ethanol for the less powerful reducing agents such as sodium triacetoxymethylborohydride and sodium cyanoborohydride. When the reducing agent is formic acid the reaction is conveniently carried out using an aqueous solution of the formic acid. The reaction is performed at a temperature in the range, for example, 10 to 100°C, such as 70 to

- 34 -

90°C or, conveniently, at or near ambient temperature. Conveniently, when the reducing agent is formic acid, protecting groups such as tert-butoxycarbonyl on the NH group to be alkylated (for example present from the synthesis of the starting material) may be removed in-situ during the reaction.

- 5 **Process (i)** For the preparation of those compounds of the Formula I wherein R<sup>1</sup> is substituted by a group T, wherein T is selected from (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (2-6C)alkanoylamino, (1-6C)alkylthio, (1-6C)alkylsulfinyl and (1-6C)alkylsulfonyl, the reaction of a compound of the Formula VII:




- 10 wherein R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, X<sup>1</sup>, n and m have any of the meanings defined hereinbefore except that any functional group is protected if necessary, R<sup>1'</sup> is a group R<sup>1</sup> as defined herein except that any T groups are replaced with Lg, and Lg is a displaceable group (for example chloro or bromo, or mesylate) with a compound of the formula TH, wherein T is as defined above except that any functional group is protected if necessary;
- 15 and thereafter any protecting group that is present is removed by conventional means.

The reaction is conveniently carried out in the presence of a suitable base. The reaction may conveniently be performed in a suitable inert solvent or diluent. Suitable bases, solvents and diluents are for example those described under process (a). The reaction is suitably performed at a temperature of for example, from 10 to 150°C, for example 30 to

20 60°C.

**Process (j)** By reacting a compound of the Formula VIII:


- 35 -



VIII

wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $X^1$ , and  $m$  have any of the meanings defined hereinbefore except that any functional group is protected if necessary and  $Lg$  is a displaceable group as hereinbefore defined,

5 with an aniline of the Formula IX:



IX

wherein  $R^5$  and  $n$  have any of the meanings defined hereinbefore except that any functional group is protected if necessary, and wherein the reaction is conveniently performed in the presence of a suitable acid,

10 and whereafter any protecting group that is present is removed by conventional means.

Suitable displaceable groups represented by  $Lg$  are as hereinbefore defined, in particular halogeno such as chloro.

The reaction is conveniently carried out in the presence of a suitable inert solvent or diluent, for example an alcohol or ester such as methanol, ethanol, isopropanol or ethyl

15 acetate, a halogenated solvent such as methylene chloride, chloroform or carbon tetrachloride, an ether such as tetrahydrofuran or 1,4-dioxane, an aromatic solvent such as toluene, or a dipolar aprotic solvent such as N,N-dimethylformamide, N,N-dimethylacetamide,

N-methylpyrrolidin-2-one acetonitrile or dimethylsulfoxide. The reaction is conveniently carried out at a temperature in the range, for example, 10 to 250°C, conveniently in the range

20 40 to 120°C or where a solvent or diluent is used at the reflux temperature. Conveniently, the compound of Formula VIII is reacted with a compound of the Formula IX in the presence of a protic solvent such as isopropanol, conveniently in the presence of an acid, for example a catalytic amount of an acid, under the conditions described above. Suitable acids include

- 36 -

hydrogen chloride gas in diethyl ether or dioxane, and hydrochloric acid, for example a 4M solution of hydrogen chloride in dioxane. Alternatively, this reaction may be conveniently carried out in an aprotic solvent, such as dioxane or a dipolar aprotic solvent such as N,N-dimethylacetamide or acetonitrile in the presence of an acid, for example hydrogen

5 chloride gas in diethyl ether or dioxane, or hydrochloric acid.


The compound of the Formula VIII, wherein Lg is halogeno, may be reacted with a compound of the Formula IX in the absence of an acid. In this reaction displacement of the halogeno leaving group Lg results in the formation of the acid HLg in-situ and auto-catalysis of the reaction. Conveniently the reaction is carried out in a suitable inert organic solvent, for

10 example isopropanol, dioxane or N,N-dimethylacetamide. Suitable conditions for this reaction are as described above.

Alternatively, the compound of Formula VIII may be reacted with a compound of the Formula IX in the presence of a suitable base. Suitable bases for this reaction are as hereinbefore defined under process (a). For example, suitable bases are alkaline earth metal

15 amides, such as sodium bis(trimethylsilyl)amide. This reaction is conveniently performed in an inert solvent or diluent, for example those mentioned above in relation to this process (j);

**Process (k)** For the preparation of those compounds of the Formula I wherein m is 1, 2 or 3, the coupling of a compound of Formula X:



X

20 where m is 1, 2 or 3 and R<sup>1</sup>, X<sup>1</sup>, R<sup>5</sup> and n are as hereinbefore defined, except any functional group is protected if necessary, with a primary or secondary amine of the formula R<sup>2</sup>NHR<sup>3</sup> where R<sup>2</sup> and R<sup>3</sup> are as defined hereinbefore; and whereafter any protecting group that is present is removed by conventional means.

The coupling reaction is conveniently carried out in the presence of a suitable coupling

25 agent, such as a carbodiimide (for example 1-[3-(dimethylamino)propyl]-

- 37 -

3-ethylcarbodiimide), or a suitable peptide coupling agent, for example

O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate (HATU).

The reaction may also be carried out in the presence of 1-hydroxybenzotriazole in conjunction with a suitable coupling agent, such as a carbodiimide. The coupling reaction is conveniently


5 carried out in an inert solvent such as, for example, a halogenated solvent such as methylene chloride, or a dipolar aprotic solvent such as N,N-dimethylformamide,

N,N-dimethylacetamide or 1-methyl-2-pyrrolidinone. Suitably the coupling reaction is carried out in the presence of a suitable base, such as an organic amine, for example

di-isopropylethylamine or 4-dimethylaminopyridine. The coupling reaction is suitably

10 performed at -25°C to 150°C, conveniently at ambient temperature.


**Process (l)** By reacting a compound of Formula IV as defined above except that any functional group is protected if necessary, with a compound of the Formula V'':



using a reductive amination procedure. Suitable reaction conditions would be apparent to the

15 skilled person and/or from the literature.

**Process (m)** For compounds of the Formula I wherein R<sup>3</sup> is (2-6C)alkyl substituted on a carbon atom by an amino, (1-6C)alkylamino, di-(1-6C)alkylamino or a saturated 5 or 6 membered heterocyclic ring which contains NR<sup>8</sup> where R<sup>8</sup> is as defined in relation to Formula I, by reacting a compound of the Formula XX:



XX

- 38 -

wherein  $R^{3a}$  is Lg-(2-6C)alkyl, wherein Lg is a displaceable group as hereinbefore defined, and wherein  $R^1$ ,  $R^2$ ,  $X^1$ ,  $R^5$ , m and n have any of the meanings defined hereinbefore except that any functional group is protected if necessary,

with ammonia or with a suitable primary or secondary amine, such as pyrrolidine,

5 and whereafter any protecting group that is present is removed by conventional means.

The reaction of process (m) is conveniently carried out in the presence of an inert solvent or diluent (for example the inert solvents and diluents described in processes (a) and (c), such as acetonitrile, N,N-dimethylacetamide, methanol, ethanol or methylene chloride). The reaction is suitably effected in the presence of a source of iodide such as sodium iodide or  
10 potassium iodide and at a temperature in the range, for example, 10 to 150°C (or the boiling point of the solvent), suitably in the range 20 to 90°C.

It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately  
15 following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution  
20 reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halogeno group.

25 Persons skilled in the art will appreciate that, in order to obtain compounds of the invention in an alternative and in some occasions, more convenient manner, the individual process steps mentioned hereinbefore may be performed in different order, and/or the individual reactions may be performed at different stage in the overall route (i.e. chemical transformations may be performed upon different intermediates to those associated  
30 hereinbefore with a particular reaction).

When a pharmaceutically-acceptable salt of a quinazoline derivative of the Formula I is required, for example an acid-addition salt, it may be obtained by, for example, reaction of said quinazoline derivative with a suitable acid using a conventional procedure. To facilitate




isolation of the compound during preparation, the compound may be prepared in the form of a salt that is not a pharmaceutically acceptable salt. The resulting salt can then be modified by conventional techniques to give a pharmaceutically acceptable salt of the compound. Such techniques include, for example ion exchange techniques or re-precipitation of the compound in the presence of a pharmaceutically acceptable counter ion. For example re-precipitation in the presence of a suitable acid such as HCl to give a hydrochloride acid addition salt.

In the section above the expression "inert solvent" refers to a solvent which does not react with the starting materials, reagents, intermediates or products in a manner which adversely affects the yield of the desired product.

10

#### Preparation of Starting Materials

Compounds of Formula II are commercially available or may be prepared using conventional techniques or analogous processes to those described in the prior art. In particular those patents and applications listed hereinbefore, such as WO 96/15118, WO 01/66099 and EP 566 226. For example, the compounds of Formula II may be prepared in accordance with Reaction Scheme 1:



Reaction Scheme 1


wherein  $R^1$ ,  $X^1$ ,  $R^5$ , Lg and n are as hereinbefore defined and Pg is a hydroxy protecting group.

- (i) Reaction suitably in an inert protic solvent (such as an alkanol for example iso-propanol), an aprotic solvent (such as dioxane) or a dipolar aprotic solvent (such as *N,N*-dimethylacetamide) in the presence of an acid, for example hydrogen chloride gas in diethyl ether or dioxane, or hydrochloric acid, under analogous conditions to those described above under process (j).

Alternatively the reaction may be carried out in one of the above inert solvents conveniently in the presence of a base, for example potassium carbonate. The above reactions are conveniently carried out at a temperature in the range, for example, 0 to 150°C, suitably at or near the reflux temperature of the reaction solvent.

- (ii) Cleavage of Pg may be performed under standard conditions for such reactions. For example when Pg is an alkanoyl group such as acetyl, it may be cleaved by heating in the presence of a methanolic ammonia solution.

- Compounds of Formula XI are known or can be prepared using known processes for the preparation of analogous compounds. If not commercially available, compounds of the Formula XI may be prepared by procedures which are selected from standard chemical techniques, techniques which are analogous to the synthesis of known, structurally similar compounds, or techniques which are analogous to the procedures described in the Examples. For example, standard chemical techniques are as described in Houben Weyl. By way of example the compound of the Formula XI in which  $R^1-X^1$  is methoxy, Lg is chloro and Pg is acetyl may be prepared using the process illustrated in Reaction Scheme 2:




Reaction Scheme 2


Reaction Scheme 2 may be generalised by the skilled man to apply to compounds within the present specification which are not specifically illustrated (for example to introduce a substituent other than methoxy at the 7-position in the quinazoline ring).

Compounds of the Formula III are commercially available or may be prepared using  
5 standard techniques, for example as illustrated in US 5,252,586 and WO 94/27965.

Compounds of the Formula IV may be prepared by reaction of a compound of Formula II with a compound of XVa or XVb:



XVa




XVb

wherein Lg is a displaceable group as hereinbefore defined and Pg is a suitable protecting  
10 group. For example, Lg may be an alkanesulfonyloxy group, such as methanesulfonyloxy and Pg may be *tert*-butylcarboxylate.

The reaction of the compound of Formula II with the compound of Formula XVa may be carried out using analogous conditions to those described in process (a) above, followed by removal of the protecting group under standard conditions. For example, the reaction may be  
15 carried out using potassium carbonate as a suitable base, *N*-methylpyrrolidin-2-one as a suitable diluent and at a temperature of about 105°C.

The reaction of the compound of Formula II with the compound of Formula XVb may be carried out under Mitsunobu conditions as described in process (e) above, followed by removal of the protecting group under standard conditions.

20 Compounds of the Formula IV may also be prepared by reaction of a compound of the Formula IX with a compound of the Formula XVc:



XVc


wherein Lg, X¹ and R¹ are as hereinbefore defined and Pg is a suitable protecting group.

The reaction of the compound of Formula IX with the compound of Formula XVc  
25 may be carried out using analogous conditions to those described in process (j) above, followed by removal of the protecting group under standard conditions.

Compounds of the Formula VI may be prepared using process (a) or process (e) above, starting with a compound prepared, for example using reaction scheme 1.

Compounds of the Formula VII may be prepared using, for example, process (a) or process (d) or process (e) in which the group represented by  $R^1$  is appropriately functionalised  
5 with a suitable displaceable group Lg such as chloro or bromo.

Compounds of the Formula VIII may be prepared using conventional methods well known in the art. For example the hydroxy protecting group, Pg, in a compound of the Formula XI as hereinbefore described in Reaction Scheme 1 is removed to give the compound of the Formula XIII:



XIII

The protecting group Pg may be removed from the compound of Formula XI using conventional techniques.

The compound of the Formula XIII may then be coupled with a compound of the  
15 Formula III as hereinbefore defined using analogous conditions to those described in process (a) or process (e).

Compounds of the Formula XX may be prepared, for example, using process (a), process (c) or process (k) above.

Compounds of the Formula V, V' and IX are commercially available or may be  
20 prepared using standard techniques. Compounds of the Formula V'' may be prepared using standard techniques, for example as illustrated in Synthesis, 1993, 12, 1233 and Tetrahedron, 1992, 48, 5557.

Compounds of the Formula X where m is 1, 2 or 3 may, for example, be prepared from a compound of the Formula IV by alkylation with  $R^1O_2C(CH_2)_m-Lg$ , wherein Lg and  $R^1$   
25 are as hereinbefore defined, in the presence of a base and using analogous conditions to those described in process (c) above, followed by transformation of the ester to the carboxylic acid (for example by saponification or acidic deprotection).

Alternatively, compounds of the Formula X where m is 1, 2 or 3 may be prepared from a compound of the Formula IV by a Mitsunobu reaction with  $R^1O_2C(CH_2)_m-OH$  using

analogous conditions to those described in process (e) above, followed by transformation of the ester to the carboxylic acid (for example by saponification or acidic deprotection).

Certain novel intermediates utilised in the above processes are provided as a further feature of the present invention together with the process for their preparation.

- 5 According to a further feature of the present invention there is provided the compounds of the Formulae VI, VII, VIII, X and XX or a salt thereof, (including pharmaceutically acceptable salts thereof), as hereinbefore defined. Examples of such compounds are 6-{{[1-(N-(2-chloroethyl)carbamoyl)piperidin-4-yl]oxy}-4-(3-chloro-2-fluoroanilino)-7-methoxyquinazoline, [4-({4-(3-chloro-2-fluoroanilino)-7-methoxyquinazolin-6-yl}oxy)piperidin-1-yl]acetic acid and [4-({4-(3-chloro-2,4-
- 10 difluoroanilino)-7-methoxyquinazolin-6-yl}oxy)piperidin-1-yl]acetic acid.

#### **Biological Assays**

- The inhibitory activities of compounds were assessed in non-cell based protein tyrosine kinase assays as well as in cell based proliferation assays before their *in vivo* activity
- 15 was assessed in Xenograft studies.

##### **a) Protein Tyrosine Kinase phosphorylation Assays**

This test measures the ability of a test compound to inhibit the phosphorylation of a tyrosine containing polypeptide substrate by EGFR or ErbB2 tyrosine kinase enzyme.

- Recombinant intracellular fragments of EGFR, erbB2 and erbB4 (accession numbers
- 20 X00588, X03363 and L07868 respectively) were cloned and expressed in the baculovirus/Sf21 system. Lysates were prepared from these cells by treatment with ice-cold lysis buffer (20mM N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES) pH7.5, 150mM NaCl, 10% glycerol, 1% Triton X-100, 1.5mM MgCl<sub>2</sub>, 1mM ethylene glycol-bis(β-aminoethyl ether) N',N',N',N'-tetraacetic acid (EGTA), plus protease inhibitors
- 25 and then cleared by centrifugation.

- Constitutive kinase activity of the recombinant protein was determined by its ability to phosphorylate a synthetic peptide (made up of a random co-polymer of Glutamic Acid, Alanine and Tyrosine in the ratio of 6:3:1). Specifically, Maxisorb™ 96-well immunoplates were coated with synthetic peptide (0.2μg of peptide in a 100μl phosphate buffered saline
- 30 (PBS) solution and incubated at 4°C overnight). Plates were washed in PBS-T (phosphate buffered saline with 0.5% Tween 20) then in 50mM HEPES pH 7.4 at room temperature to remove any excess unbound synthetic peptide. EGFR, ErbB2 or ErbB4 tyrosine kinase activity was assessed by incubation in peptide coated plates for 20 minutes at 22°C in 100mM

- 44 -

HEPES pH 7.4, adenosine triphosphate (ATP) at  $K_m$  concentration for the respective enzyme, 10mM  $MnCl_2$ , 0.1mM  $Na_3VO_4$ , 0.2mM DL-dithiothreitol (DTT), 0.1% Triton X-100 with test compound in DMSO (final concentration of 2.5%). Reactions were terminated by the removal of the liquid components of the assay followed by washing of the plates with

5 PBS-T.

The immobilised phospho-peptide product of the reaction was detected by immunological methods. Firstly, plates were incubated for 90 minutes at room temperature with anti-phosphotyrosine primary antibodies that were raised in the mouse (4G10 from Upstate Biotechnology). Following extensive washing, plates were treated with Horseradish  
10 Peroxidase (HRP) conjugated sheep anti-mouse secondary antibody (NXA931 from Amersham) for 60 minutes at room temperature. After further washing, HRP activity in each well of the plate was measured colorimetrically using 22'-Azino-di-[3-ethylbenzthiazoline sulfonate (6)] diammonium salt crystals (ABTS™ from Roche) as a substrate.

Quantification of colour development and thus enzyme activity was achieved by the  
15 measurement of absorbance at 405nm on a Molecular Devices ThermoMax microplate reader. Kinase inhibition for a given compound was expressed as an  $IC_{50}$  value. This was determined by calculation of the concentration of compound that was required to give 50% inhibition of phosphorylation in this assay. The range of phosphorylation was calculated from the positive (vehicle plus ATP) and negative (vehicle minus ATP) control values.

20 **b) EGFR driven KB cell proliferation assay**

This assay measures the ability of a test compound to inhibit the proliferation of KB cells (human naso-pharyngeal carcinoma obtained from the American Type Culture Collection (ATCC).

KB cells (human naso-pharyngeal carcinoma obtained from the ATCC were cultured  
25 in Dulbecco's modified Eagle's medium (DMEM) containing 10% foetal calf serum, 2 mM glutamine and non-essential amino acids at 37°C in a 7.5%  $CO_2$  air incubator. Cells were harvested from the stock flasks using Trypsin/ethylaminediaminetetraacetic acid (EDTA). Cell density was measured using a haemocytometer and viability was calculated using trypan blue solution before being seeded at a density of  $1.25 \times 10^3$  cells per well of a 96 well plate in  
30 DMEM containing 2.5% charcoal stripped serum, 1mM glutamine and non-essential amino acids at 37°C in 7.5%  $CO_2$  and allowed to settle for 4 hours.

Following adhesion to the plate, the cells are treated with or without EGF (final concentration of 1ng/ml) and with or without compound at a range of concentrations in

- 45 -

dimethylsulfoxide (DMSO) (0.1% final) before incubation for 4 days. Following the incubation period, cell numbers were determined by addition of 50µl of 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (stock 5mg/ml) for 2 hours. MTT solution was then tipped off, the plate gently tapped dry and the cells dissolved upon the addition of 100µl of DMSO.

Absorbance of the solubilised cells was read at 540nm using a Molecular Devices ThermoMax microplate reader. Inhibition of proliferation was expressed as an IC<sub>50</sub> value. This was determined by calculation of the concentration of compound that was required to give 50% inhibition of proliferation. The range of proliferation was calculated from the positive (vehicle plus EGF) and negative (vehicle minus EGF) control values.

**c) *In vivo* Xenograft assays**

**(i) LOVO**

This assay measures the ability of a test compound to inhibit the growth of a LoVo tumour (colorectal adenocarcinoma obtained from the ATCC) in Female Swiss athymic mice (Alderley Park, *nu/nu* genotype).

Female Swiss athymic (*nu/nu* genotype) mice were bred and maintained in Alderley Park in negative pressure Isolators (PFI Systems Ltd.). Mice were housed in a barrier facility with 12hr light/dark cycles and provided with sterilised food and water *ad libitum*. All procedures were performed on mice of at least 8 weeks of age. LoVo tumour cell (colorectal adenocarcinoma obtained from the ATCC) xenografts were established in the hind flank of donor mice by sub cutaneous injections of  $1 \times 10^7$  freshly cultured cells in 100µl of serum free media per animal. On day 5 post-implant, mice were randomised into groups of 7 prior to the treatment with compound or vehicle control that was administered once daily at 0.1ml/10g body weight. Tumour volume was assessed twice weekly by bilateral Vernier calliper measurement, using the formula  $(\text{length} \times \text{width}) \times \sqrt{(\text{length} \times \text{width}) \times (\pi/6)}$ , where length was the longest diameter across the tumour, and width was the corresponding perpendicular. Growth inhibition from start of study was calculated by comparison of the mean changes in tumour volume for the control and treated groups, and statistical significance between the two groups was evaluated using a Students *t* test.

**(ii) *In vivo* BT-474 Xenograft assay**

This assay measures the ability of a test compound to inhibit the growth of a BT-474 tumour cell xenograft (human mammary carcinoma obtained from Dr Baselga, Laboratorio Recerca Oncologica, Paseo Vall D'Hebron 119-129, Barcelona 08035, Spain) in Female

Swiss athymic mice (Alderley Park, *nu/nu* genotype) (Baselga, J. *et al.* (1998) *Cancer Research*, **58**, 2825-2831).

Female Swiss athymic (*nu/nu* genotype) mice were bred and maintained in Alderley Park in negative pressure Isolators (PFI Systems Ltd.). Mice were housed in a barrier facility with 12hr light/dark cycles and provided with sterilised food and water *ad libitum*. All procedures were performed on mice of at least 8 weeks of age. BT-474 tumour cell xenografts were established in the hind flank of donor mice by sub-cutaneous injections of  $1 \times 10^7$  freshly cultured cells in 100  $\mu$ l of serum free media with 50% Matrigel per animal. On day 14 post-implant, mice were randomised into groups of 10 prior to the treatment with compound or vehicle control that was administered once daily at 0.1ml/kg body weight. Tumour volume was assessed twice weekly by bilateral Vernier calliper measurement, using the formula  $(\text{length} \times \text{width}) \times \sqrt{(\text{length} \times \text{width}) \times (\pi/6)}$ , where length was the longest diameter across the tumour, and width was the corresponding perpendicular. Growth inhibition from start of treatment was calculated by comparison of the mean changes in tumour volume for the control and treated groups, and statistical significance between the two groups was evaluated using a Students *t* test.

#### d) hERG-encoded Potassium Channel Inhibition Assay

This assay determines the ability of a test compound to inhibit the tail current flowing through the human ether-a-go-go-related-gene (hERG)-encoded potassium channel.

Human embryonic kidney (HEK) cells expressing the hERG-encoded channel were grown in Minimum Essential Medium Eagle (EMEM; Sigma-Aldrich catalogue number M2279), supplemented with 10% Foetal Calf Serum (Labtech International; product number 4-101-500), 10% M1 serum-free supplement (Egg Technologies; product number 70916) and 0.4 mg/ml Geneticin G418 (Sigma-Aldrich; catalogue number G7034). One or two days before each experiment, the cells were detached from the tissue culture flasks with Accutase (TCS Biologicals) using standard tissue culture methods. They were then put onto glass coverslips resting in wells of a 12 well plate and covered with 2 ml of the growing media.

For each cell recorded, a glass coverslip containing the cells was placed at the bottom of a Perspex chamber containing bath solution (see below) at room temperature ( $\sim 20^\circ\text{C}$ ).

This chamber was fixed to the stage of an inverted, phase-contrast microscope. Immediately after placing the coverslip in the chamber, bath solution was perfused into the chamber from a gravity-fed reservoir for 2 minutes at a rate of  $\sim 2$  ml/min. After this time, perfusion was stopped.



- 47 -

A patch pipette made from borosilicate glass tubing (GC120F, Harvard Apparatus) using a P-97 micropipette puller (Sutter Instrument Co.) was filled with pipette solution (see hereinafter). The pipette was connected to the headstage of the patch clamp amplifier (Axopatch 200B, Axon Instruments) via a silver/silver chloride wire. The headstage ground was connected to the earth electrode. This consisted of a silver/silver chloride wire embedded in 3% agar made up with 0.85% sodium chloride.

The cell was recorded in the whole cell configuration of the patch clamp technique. Following "break-in", which was done at a holding potential of  $-80$  mV (set by the amplifier), and appropriate adjustment of series resistance and capacitance controls, electrophysiology software (*Clampex*, Axon Instruments) was used to set a holding potential ( $-80$  mV) and to deliver a voltage protocol. This protocol was applied every 15 seconds and consisted of a 1 s step to  $+40$  mV followed by a 1 s step to  $-50$  mV.

The current response to each imposed voltage protocol was low pass filtered by the amplifier at 1 kHz. The filtered signal was then acquired, on line, by digitising this analogue signal from the amplifier with an analogue to digital converter. The digitised signal was then captured on a computer running *Clampex* software (Axon Instruments). During the holding potential and the step to  $+40$  mV the current was sampled at 1 kHz. The sampling rate was then set to 5 kHz for the remainder of the voltage protocol.

The compositions, pH and osmolarity of the bath and pipette solution are tabulated below.

Salt	Pipette (mM)	Bath (mM)
NaCl	-	137
KCl	130	4
MgCl <sub>2</sub>	1	1
CaCl <sub>2</sub>	-	1.8
HEPES	10	10
glucose	-	10
Na <sub>2</sub> ATP	5	-
EGTA	5	-

Parameter	Pipette	Bath
pH	7.18 – 7.22	7.40
pH adjustment with	1M KOH	1M NaOH
Osmolarity (mOsm)	275-285	285-295

- 48 -

The amplitude of the hERG-encoded potassium channel tail current following the step from +40 mV to -50 mV was recorded on-line by *Clampex* software (Axon Instruments). Following stabilisation of the tail current amplitude, bath solution containing the vehicle for the test substance was applied to the cell. Providing the vehicle application had no significant effect on tail current amplitude, a cumulative concentration effect curve to the compound was then constructed.

The effect of each concentration of test compound was quantified by expressing the tail current amplitude in the presence of a given concentration of test compound as a percentage of that in the presence of vehicle.

10 Test compound potency ( $IC_{50}$ ) was determined by fitting the percentage inhibition values making up the concentration-effect to a four parameter Hill equation using a standard data-fitting package. If the level of inhibition seen at the highest test concentration did not exceed 50%, no potency value was produced and a percentage inhibition value at that concentration was quoted.

15 e) Clone 24 phospho-erbB2 cell assay

This immunofluorescence end point assay measures the ability of a test compound to inhibit the phosphorylation of erbB2 in a MCF7 (breast carcinoma) derived cell line which was generated by transfecting MCF7 cells with the full length erbB2 gene using standard methods to give a cell line that overexpresses full length wild type erbB2 protein (hereinafter 20 'Clone 24' cells).

Clone 24 cells were cultured in Growth Medium (phenol red free Dulbecco's modified Eagle's medium (DMEM) containing 10% foetal bovine serum, 2 mM glutamine and 1.2mg/ml G418) in a 7.5% CO<sub>2</sub> air incubator at 37°C. Cells were harvested from T75 stock flasks by washing once in PBS (phosphate buffered saline, pH7.4, Gibco No. 25 10010-015) and harvested using 2mls of Trypsin (1.25mg/ml) / ethylenediaminetetraacetic acid (EDTA) (0.8mg/ml) solution. The cells were resuspended in Growth Medium. Cell density was measured using a haemocytometer and viability was calculated using Trypan Blue solution before being further diluted in Growth Medium and seeded at a density of  $1 \times 10^4$  cells per well (in 100ul) into clear bottomed 96 well plates (Packard, No. 6005182).

30 3 days later, Growth Medium was removed from the wells and replaced with 100ul Assay Medium (phenol red free DMEM, 2mM glutamine, 1.2mg/ml G418) either with or without erbB inhibitor compound. Plates were returned to the incubator for 4hrs and then 20ul of 20% formaldehyde solution in PBS was added to each well and the plate was left at room

- 49 -

temperature for 30 minutes. This fixative solution was removed with a multichannel pipette, 100µl of PBS was added to each well and then removed with a multichannel pipette and then 50µl PBS was added to each well. Plates were then sealed and stored for up to 2 weeks at 4°C.

Immunostaining was performed at room temperature. Wells were washed once with 5 200µl PBS / Tween 20 (made by adding 1 sachet of PBS / Tween dry powder (Sigma, No. P3563) to 1L of double distilled H<sub>2</sub>O) using a plate washer then 200µl Blocking Solution (5% Marvel dried skimmed milk (Nestle) in PBS / Tween 20) was added and incubated for 10 minutes. Blocking Solution was removed using a plate washer and 200µl of 0.5% Triton X-100 / PBS was added to permeabilise the cells. After 10 minutes, the plate was washed 10 with 200µl PBS / Tween 20 and then 200µl Blocking Solution was added once again and incubated for 15 minutes. Following removal of the Blocking Solution with a plate washer, 30µl of rabbit polyclonal anti-phospho ErbB2 IgG antibody (epitope phospho-Tyr 1248, SantaCruz, No. SC-12352-R), diluted 1:250 in Blocking Solution, was added to each well and incubated for 2 hours. Then this primary antibody solution was removed from the wells using 15 a plate washer followed by two 200µl PBS / Tween 20 washes using a plate washer. Then 30µl of Alexa-Fluor 488 goat anti-rabbit IgG secondary antibody (Molecular Probes, No. A-11008), diluted 1:750 in Blocking Solution, was added to each well. From now onwards, wherever possible, plates were protected from light exposure, at this stage by sealing with black backing tape. The plates were incubated for 45 minutes and then the secondary 20 antibody solution was removed from the wells followed by two 200µl PBS / Tween 20 washes using a plate washer. Then 100µl PBS was added to each plate, incubated for 10 minutes and then removed using a plate washer. Then a further 100µl PBS was added to each plate and then, without prolonged incubation, removed using a plate washer. Then 50µl of PBS was added to each well and plates were resealed with black backing tape and stored for up to 2 25 days at 4°C before analysis.

The Fluorescence signal in each well was measured using an Acumen Explorer Instrument (Acumen Bioscience Ltd.), a plate reader that can be used to rapidly quantitate features of images generated by laser-scanning. The instrument was set to measure the number of fluorescent objects above a pre-set threshold value and this provided a measure of 30 the phosphorylation status of erbB2 protein. Fluorescence dose response data obtained with each compound was exported into a suitable software package (such as Origin) to perform curve fitting analysis. Inhibition of erbB2 phosphorylation was expressed as an IC<sub>50</sub> value.

This was determined by calculation of the concentration of compound that was required to give 50% inhibition of erbB2 phosphorylation signal.

Although the pharmacological properties of the compounds of the Formula I vary with structural change as expected, in general activity possessed by compounds of the Formula I, 5 may be demonstrated at the following concentrations or doses in one or more of the above tests:-

- Test (a):-  $IC_{50}$  in the range, for example, 0.001 - 10  $\mu M$ ;
- Test (b):-  $IC_{50}$  in the range, for example, 0.001 - 10  $\mu M$ ;
- Test (e):-  $IC_{50}$  in the range, for example, 0.001 - 10  $\mu M$ ;
- 10 Test (c):- activity in the range, for example, 1-200 mg/kg/day;

By way of example, Table A illustrates the activity of representative compounds according to the invention. Column 2 of Table A shows  $IC_{50}$  data from Test (a) for the inhibition of EGFR tyrosine kinase protein phosphorylation; column 3 shows  $IC_{50}$  data from Test (a) for the inhibition of erbB2 tyrosine kinase protein phosphorylation; column 4 shows 15  $IC_{50}$  data for inhibition of proliferation of KB cells in Test (b) described above; and column 5 shows  $IC_{50}$  data for inhibition of phosphorylation of erbB2 in a MCF7 derived cell line in Test (e) described above:

Table A

Example Number	$IC_{50}$ ( $\mu M$ ) Test (a): Inhibition of EGFR tyrosine kinase protein phosphorylation	$IC_{50}$ ( $\mu M$ ) Test (a): Inhibition of erbB2 tyrosine kinase protein phosphorylation	$IC_{50}$ ( $\mu M$ ) Test (b): EGFR driven KB cell proliferation assay	$IC_{50}$ ( $\mu M$ ) Test (e): Inhibition of erbB2 tyrosine kinase protein phosphorylation
5	0.004	0.047	0.009	0.006
7	0.003	0.013	0.017	0.014
10	0.004	0.010	-	0.013

20 According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier.

The compositions of the invention may be in a form suitable for oral use (for example

- 51 -

as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intramuscular dosing or as a suppository for rectal dosing).

The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and the particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 0.5 g of active agent (more suitably from 0.5 to 100 mg, for example from 1 to 30 mg) compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition.

The size of the dose for therapeutic or prophylactic purposes of a compound of the Formula I will naturally vary according to the nature and severity of the conditions, the age and sex of the animal or patient and the route of administration, according to well known principles of medicine.

In using a compound of the Formula I for therapeutic or prophylactic purposes it will generally be administered so that a daily dose in the range, for example, 0.1 mg/kg to 75 mg/kg body weight is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed. Thus, for example, for intravenous administration, a dose in the range, for example, 0.1 mg/kg to 30 mg/kg body weight will generally be used. Similarly, for administration by inhalation, a dose in the range, for example, 0.05 mg/kg to 25 mg/kg body weight will be used. Oral administration is however preferred, particularly in tablet form. Typically, unit dosage forms will contain about 0.5 mg to 0.5 g of a compound of this invention.

- 52 -

We have found that the compounds of the present invention possess anti-proliferative properties such as anti-cancer properties that are believed to arise from their erbB family receptor tyrosine kinase inhibitory activity, and particularly a mixed erbB2/ EGF profile.

Accordingly, the compounds of the present invention are expected to be useful in the treatment of diseases or medical conditions mediated alone or in part by erbB receptor tyrosine kinases, i.e. the compounds may be used to produce an erbB receptor tyrosine kinase inhibitory effect in a warm-blooded animal in need of such treatment. Thus the compounds of the present invention provide a method for the treatment of malignant cells characterised by inhibition of one or more of the erbB family of receptor tyrosine kinases. Particularly the compounds of the invention may be used to produce an anti-proliferative and/or pro-apoptotic and/or anti-invasive effect mediated alone or in part by the inhibition of erbB receptor tyrosine kinases. Particularly, the compounds of the present invention are expected to be useful in the prevention or treatment of those tumours that are sensitive to inhibition of one or more of the erbB receptor tyrosine kinases, that are involved in the signal transduction steps which drive proliferation and survival of these tumour cells. Accordingly the compounds of the present invention are expected to be useful in the treatment of psoriasis, benign prostatic hyperplasia (BPH), atherosclerosis and restenosis and/or cancer by providing an anti-proliferative effect, particularly in the treatment of erbB receptor tyrosine kinase sensitive cancers. Such benign or malignant tumours may affect any tissue and include non-solid tumours such as leukaemia, multiple myeloma or lymphoma, and also solid tumours, for example bile duct, bone, bladder, brain/CNS, breast, colorectal, endometrial, gastric, head and neck, hepatic, lung, neuronal, oesophageal, ovarian, pancreatic, prostate, renal, skin, testicular, thyroid, uterine and vulval cancers.

According to this aspect of the invention there is provided a compound of the Formula I, or a pharmaceutically acceptable salt thereof, for use as a medicament.

According to a further aspect of the invention there is provided a compound of the Formula I, or a pharmaceutically acceptable salt thereof, for use in the production of an anti-proliferative effect in a warm-blooded animal such as man.

Thus according to this aspect of the invention there is provided the use of a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of an anti-proliferative effect in a warm-blooded animal such as man.

- 53 -

According to a further feature of this aspect of the invention there is provided a method for producing an anti-proliferative effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a quinazoline derivative of the Formula I, or a pharmaceutically acceptable salt thereof, as  
5 hereinbefore defined.

According to a further aspect of the invention there is provided the use of a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the prevention or treatment of those tumours which are sensitive to inhibition of erbB receptor tyrosine kinases,  
10 such as a combination of EGFR and erbB2, that are involved in the signal transduction steps which lead to the proliferation of tumour cells.

According to a further feature of this aspect of the invention there is provided a method for the prevention or treatment of those tumours which are sensitive to inhibition of one or more of the erbB family of receptor tyrosine kinases, such as a combination of EGFR  
15 and erbB2, that are involved in the signal transduction steps which lead to the proliferation and/or survival of tumour cells which comprises administering to said animal an effective amount of a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore.

According to a further feature of this aspect of the invention there is provided a  
20 compound of the Formula I, or a pharmaceutically acceptable salt thereof, for use in the prevention or treatment of those tumours which are sensitive to inhibition of erbB receptor tyrosine kinases, such as a combination of EGFR and erbB2, that are involved in the signal transduction steps which lead to the proliferation of tumour cells.

According to a further aspect of the invention there is provided the use of a  
25 quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in providing a combined EGFR and erbB2 tyrosine kinase inhibitory effect.

According to a further feature of this aspect of the invention there is provided a method for providing a combined EGFR and erbB2 tyrosine kinase inhibitory effect which  
30 comprises administering to said animal an effective amount of a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore.

- 54 -

According to a further feature of this aspect of the invention there is provided a compound of the Formula I, or a pharmaceutically acceptable salt thereof, for use in providing a combined EGFR and erbB2 tyrosine kinase inhibitory effect.

According to a further aspect of the present invention there is provided the use of a  
5 quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the treatment of a cancer (for example a cancer selected from leukaemia, multiple myeloma, lymphoma, bile duct, bone, bladder, brain/CNS, breast, colorectal, endometrial, gastric, head and neck, hepatic, lung, neuronal, oesophageal, ovarian, pancreatic, prostate, renal, skin, testicular, thyroid,  
10 uterine and vulval cancer).

According to a further feature of this aspect of the invention there is provided a method for treating a cancer (for example a cancer selected from leukaemia, multiple myeloma, lymphoma, bile duct, bone, bladder, brain/CNS, breast, colorectal, endometrial, gastric, head and neck, hepatic, lung, neuronal, oesophageal, ovarian, pancreatic, prostate,  
15 renal, skin, testicular, thyroid, uterine and vulval cancer) in a warm-blooded animal, such as man, in need of such treatment, which comprises administering to said animal an effective amount of a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore.

According to a further aspect of the invention there is provided a compound of the  
20 Formula I, or a pharmaceutically acceptable salt thereof, for use in the treatment of a cancer (for example selected from leukaemia, multiple myeloma, lymphoma, bile duct, bone, bladder, brain/CNS, breast, colorectal, endometrial, gastric, head and neck, hepatic, lung, neuronal, oesophageal, ovarian, pancreatic, prostate, renal, skin, testicular, thyroid, uterine and vulval cancer).

25 As mentioned above the size of the dose required for the therapeutic or prophylactic treatment of a particular disease will necessarily be varied depending upon, amongst other things, the host treated, the route of administration and the severity of the illness being treated.

The anti-proliferative treatment defined hereinbefore may be applied as a sole therapy  
30 or may involve, in addition to the quinazoline derivative of the invention, conventional surgery or radiotherapy or chemotherapy. Such chemotherapy may include one or more of the following categories of anti-tumour agents:-



- 55 -

- (i) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology, such as alkylating agents (for example cis-platin, carboplatin, cyclophosphamide, nitrogen mustard, melphalan, chlorambucil, busulphan and nitrosoureas); antimetabolites (for example antifolates such as fluoropyrimidines like 5-fluorouracil and tegafur, raltitrexed, 5 methotrexate, cytosine arabinoside and hydroxyurea; antitumour antibiotics (for example anthracyclines like adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin and mithramycin); antimitotic agents (for example vinca alkaloids like vincristine, vinblastine, vindesine and vinorelbine and taxoids like taxol and taxotere); and topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and 10 teniposide, amsacrine, topotecan and camptothecin);
- (ii) cytostatic agents such as antioestrogens (for example tamoxifen, toremifene, raloxifene, droloxifene and idoxifene), oestrogen receptor down regulators (for example fulvestrant), antiandrogens (for example bicalutamide, flutamide, nilutamide and cyproterone acetate), LHRH antagonists or LHRH agonists (for example goserelin, leuprorelin and 15 buserelin), progestogens (for example megestrol acetate), aromatase inhibitors (for example as anastrozole, letrozole, vorazole and exemestane) and inhibitors of 5 $\alpha$ -reductase such as finasteride;
- (iii) agents which inhibit cancer cell invasion (for example metalloproteinase inhibitors like marimastat and inhibitors of urokinase plasminogen activator receptor function);
- 20 (iv) inhibitors of growth factor function, for example such inhibitors include growth factor antibodies, growth factor receptor antibodies (for example the anti-erbB2 antibody trastuzumab [Herceptin<sup>TM</sup>] and the anti-erbB1 antibody cetuximab [C225]), farnesyl transferase inhibitors, tyrosine kinase inhibitors and serine/threonine kinase inhibitors, for example other inhibitors of the epidermal growth factor family (for example EGFR family 25 tyrosine kinase inhibitors such as
- N-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine (gefitinib, AZD1839), N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI-774) and 6-acrylamido-N-(3-chloro-4-fluorophenyl)-7-(3-morpholinopropoxy)quinazolin-4-amine (CI 30 1033)), for example inhibitors of the platelet-derived growth factor family and for example inhibitors of the hepatocyte growth factor family;
- (v) antiangiogenic agents such as those which inhibit the effects of vascular endothelial growth factor, (for example the anti-vascular endothelial cell growth factor antibody

- 56 -

bevacizumab [Avastin™], compounds such as those disclosed in International Patent Applications WO 97/22596, WO 97/30035, WO 97/32856 and WO 98/13354) and compounds that work by other mechanisms (for example linomide, inhibitors of integrin  $\alpha v \beta 3$  function and angiostatin);

- 5 (vi) vascular damaging agents such as Combretastatin A4 and compounds disclosed in International Patent Applications WO 99/02166, WO00/40529, WO 00/41669, WO01/92224, WO02/04434 and WO02/08213;
- (vii) antisense therapies, for example those which are directed to the targets listed above, such as ISIS 2503, an anti-ras antisense;
- 10 (viii) gene therapy approaches, including for example approaches to replace aberrant genes such as aberrant p53 or aberrant BRCA1 or BRCA2, GDEPT (gene-directed enzyme pro-drug therapy) approaches such as those using cytosine deaminase, thymidine kinase or a bacterial nitroreductase enzyme and approaches to increase patient tolerance to chemotherapy or radiotherapy such as multi-drug resistance gene therapy; and
- 15 (ix) immunotherapy approaches, including for example ex-vivo and in-vivo approaches to increase the immunogenicity of patient tumour cells, such as transfection with cytokines such as interleukin 2, interleukin 4 or granulocyte-macrophage colony stimulating factor, approaches to decrease T-cell anergy, approaches using transfected immune cells such as cytokine-transfected dendritic cells, approaches using cytokine-transfected tumour cell lines
- 20 and approaches using anti-idiotypic antibodies.

Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate dosing of the individual components of the treatment. Such combination products employ the compounds of this invention within the dosage range described hereinbefore and the other pharmaceutically-active agent within its approved dosage range.

- 25 According to this aspect of the invention there is provided a pharmaceutical product comprising a quinazoline derivative of the Formula I as defined hereinbefore and an additional anti-tumour agent as defined hereinbefore for the conjoint treatment of cancer.

Although the compounds of the Formula I are primarily of value as therapeutic agents for use in warm-blooded animals (including man), they are also useful whenever it is required

30 to inhibit the effects of the erbB receptor tyrosine protein kinases. Thus, they are useful as pharmacological standards for use in the development of new biological tests and in the search for new pharmacological agents.

- 57 -

The invention will now be illustrated by the following non-limiting examples in which, unless stated otherwise:

- (i) temperatures are given in degrees Celsius (°C); operations were carried out at room or ambient temperature, that is, at a temperature in the range of 18-25°C;
- 5 (ii) organic solutions were dried over anhydrous magnesium sulfate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 Pascals; 4.5-30mmHg) with a bath temperature of up to 60°C;
- (iii) chromatography means flash chromatography on silica gel; thin layer chromatography (TLC) was carried out on silica gel plates;
- 10 (iv) in general, the course of reactions was followed by TLC and / or analytical LCMS, and reaction times are given for illustration only;
- (v) final products had satisfactory proton nuclear magnetic resonance (NMR) spectra and/or mass spectral data;
- (vi) yields are given for illustration only and are not necessarily those which can be obtained
- 15 by diligent process development; preparations were repeated if more material was required;
- (vii) when given, NMR data is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard, determined at 400 MHz using perdeuterio dimethyl sulfoxide (DMSO-d<sub>6</sub>) as solvent unless otherwise indicated; the following abbreviations have been used: s, singlet; d, doublet; t,
- 20 triplet; q, quartet; m, multiplet; b, broad;
- (viii) chemical symbols have their usual meanings; SI units and symbols are used;
- (ix) solvent ratios are given in volume:volume (v/v) terms; and
- (x) mass spectra (MS) were run with an electron energy of 70 electron volts in the chemical ionization (CI) mode using a direct exposure probe and ionization was effected by
- 25 electrospray; values for m/z are given; generally, only ions which indicate the parent mass are reported; and unless otherwise stated, the mass ion quoted is (MH)<sup>+</sup>;
- (xi) unless stated otherwise compounds containing an asymmetrically substituted carbon and/or sulfur atom have not been resolved;
- (xii) where a synthesis is described as being analogous to that described in a previous example
- 30 the amounts used are the millimolar ratio equivalents to those used in the previous example;
- (xiii) the following abbreviations have been used:

DCM                      dichloromethane;

DMF                      *N,N*-dimethylformamide;

- 58 -

DMA *N,N*-dimethylacetamide;

THF Tetrahydrofuran;

(xiv) where a synthesis is described as leading to an acid addition salt (e.g. HCl salt), the specific stoichiometry of the salt was not confirmed.

- 5 (xv) In Examples 1 to 12 unless otherwise stated, all NMR data is reported on free-base material, with isolated salts converted to the free-base form prior to characterisation.

### Example 1

#### Preparation of 4-(3-Chloro-2-fluoroanilino)-7-methoxy-6-[[1-(N-methylcarbamoylmethyl)piperidin-4-yl]oxy]quinazoline

- 10 2-Chloro-N-methylacetamide (32 mg, 0.3 mmol) was added dropwise to a mixture of 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-[(piperidin-4-yl)oxy]quinazoline (120 mg, 0.3 mmol), potassium iodide (16 mg, 0.1 mmol), and potassium carbonate (50 mg, 0.36 mmol) in acetonitrile (5 ml). The mixture was heated at reflux for one hour. After evaporation of the solvents under vacuum, the residue was taken up in dichloromethane. The organic solution  
15 was washed with water and brine, dried over magnesium sulfate. After evaporation of the solvents under vacuum, the residue was purified by chromatography on silica gel (eluant: 1% to 2% 7N methanolic ammonia in dichloromethane) to give the title compound as a white solid (85 mg, 60%).

- <sup>1</sup>H NMR Spectrum: (CDCl<sub>3</sub>) 1.98 (m, 2H), 2.08 (m, 2H), 2.46 (m, 2H), 2.85 (m, 2H),  
20 2.87 (d, 3H), 3.07 (s, 2H), 4.02 (s, 3H), 4.49 (m, 1H), 7.16 (m, 4H), 7.31 (m, 2H), 8.49 (m, 1H), 8.71 (s, 1H); Mass spectrum: MH<sup>+</sup> 474

4-(3-Chloro-2-fluoroanilino)-7-methoxy-6-[(piperidin-4-yl)oxy]quinazoline used as the starting material was prepared as follows:

### Step 1

- 25 **6-Acetoxy-4-(3-chloro-2-fluoroanilino)-7-methoxyquinazoline hydrochloride**

- 6-Acetoxy-4-chloro-7-methoxyquinazoline (prepared as described in Example 25-5 of  
in WO01/66099, 6.00 g, 23.8 mmol) and 3-chloro-2-fluoroaniline (3.46 g, 23.8 mmol) were  
suspended in *iso*-propanol (200 ml). The mixture was heated to 80°C under reflux for 3  
hours. The solvent was evaporated; the residue was crystallised from acetonitrile, giving the  
30 product hydrochloride as a pale pink crystalline solid (8.16 g, 92%);

<sup>1</sup>H NMR: 2.37 (s, 3H), 4.00 (s, 3H), 7.34 (ddd, 1H), 7.48 (s, 1H), 7.52 (ddd, 1H), 7.61  
(ddd, 1H), 8.62 (s, 1H), 8.86 (s, 1H); Mass Spectrum: 362.4, 364.4.

- 59 -

**Step 2****4-(3-Chloro-2-fluoroanilino)-6-hydroxy-7-methoxyquinazoline**

6-Acetoxy-4-(3-chloro-2-fluoroanilino)-7-methoxyquinazoline hydrochloride from step 1 (8.72 g, 21.9 mmol) was dissolved in methanol (200 ml). Concentrated aqueous ammonia (15 ml) was added, and the solution heated to 50°C with stirring for 2 hours, causing precipitation of a cream coloured solid. The solid was collected by filtration, washed with diethyl ether (3x 200 ml), and dried *in vacuo* at 60°C over diphosphorous pentoxide, giving the product as an off white solid (5.40 g, 77%);

<sup>1</sup>H NMR: 3.95 (s, 3H), 7.19 (s, 1H), 7.23 (dd, 1H), 7.42 (dd, 1H), 7.50 (dd, 1H), 7.64 (s, 1H), 8.32 (s, 1H), 9.43 (s, 1H), 9.67 (br.s, 1H); Mass Spectrum: 320.4, 322.4.

**Step 3****6-([(1-*tert*-Butoxycarbonyl)piperidin-4-yl]oxy)-4-(3-chloro-2-fluoroanilino)-7-methoxyquinazoline**

4-(3-Chloro-2-fluoroanilino)-6-hydroxy-7-methoxyquinazoline from Step 2 (1870 mg, 5.85 mmol) was dissolved in DMA (50 ml). *tert*-Butyl (4-methanesulfonyloxy)piperidine-1-carboxylate (prepared as in Chemical & Pharmaceutical Bulletin 2001, 49(7), 822-829; 490 mg, 1.76 mmol) and cesium fluoride (890 mg, 5.85 mmol) were added, and the mixture was heated to 85°C with stirring. At intervals of 2 hours, 4 hours and 6 hours, *tert*-butyl 4-methanesulfonyloxypiperidine-1-carboxylate and cesium fluoride were added in the above quantities to the reaction mixture. Heating was continued at 85°C for a further 6 hours after the final addition. The solvent was evaporated, and the residue was partitioned between DCM (150 ml) and H<sub>2</sub>O (150 ml). The aqueous layer was extracted with DCM (4x 100 ml), and the extractions combined with the DCM layer. The combined DCM fractions were dried over MgSO<sub>4</sub> and evaporated. The residue was purified by chromatography, eluting with 0 to 2.5% (7:1 MeOH / concentrated aqueous NH<sub>4</sub>OH) in DCM. The appropriate fractions were combined and evaporated, giving the product as a light brown foam (2.40 g, 58%, allowing for 2.3 equivalents of residual DMA);

<sup>1</sup>H NMR: 1.40 (s, 9H), 1.60-1.65 (m, 2H), 1.95-2.00 (m, 2H), 3.20-3.25 (m, 2H), 3.65-3.70 (m, 2H), 3.92 (s, 3H), 4.68 (m, 1H), 7.21 (s, 1H), 7.27 (dd, 1H), 7.47 (ddd, 1H), 7.51 (dd, 1H), 7.85 (s, 1H), 8.36 (s, 1H), 9.53 (s, 1H); Mass Spectrum: 503.5, 505.5.

**Step 4****4-(3-chloro-2-fluoroanilino)-7-methoxy-6-[(piperidin-4-yl)oxy]quinazoline**

6-[[1-*tert*-Butoxycarbonyl]piperidin-4-yl]oxy}-4-(3-chloro-2-fluoroanilino)-7-methoxyquinazoline from step 3 (350 mg, 0.70 mmol) was dissolved in trifluoroacetic acid (5 ml), and the solution stood for 2 hours. The excess trifluoroacetic acid was evaporated, and the residue was azeotroped twice with DCM. The residue was purified by chromatography, eluting with 0 to 4% (7:1 MeOH / concentrated aqueous NH<sub>4</sub>OH) in DCM. Evaporation of the appropriate fractions gave the product as an off-white solid (270 mg, 96%);

<sup>1</sup>H NMR: 1.53-1.64 (m, 2H), 2.00-2.05 (m, 2H), 2.64-2.72 (m, 2H), 3.00-3.07 (m, 2H), 3.92 (s, 3H), 4.60 (m, 1H), 7.20 (s, 1H), 7.26 (ddd, 1H), 7.47 (dd, 1H), 7.50 (dd, 1H), 7.82 (s, 1H), 8.34 (s, 1H), 9.56 (s, 1H); Mass Spectrum: 403.2, 405.2.

**Example 2****Preparation of 4-(3-Chloro-2-fluoroanilino)-7-methoxy-6-[[1-(N-methylcarbamoyl)piperidin-4-yl]oxy] quinazoline**

Methylisocyanate (20.4 µl, 0.33 mmol) was added dropwise to a mixture of 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-[(piperidin-4-yl)oxy]quinazoline (120 mg, 0.3 mmol) in dichloromethane (5 ml) at room temperature. The mixture was stirred at room temperature for 4 hours. After evaporation of the solvents under vacuum, the residue was purified by chromatography on silica gel (eluant: 2% 7N methanolic ammonia in dichloromethane) to give the title compound as a white solid (100 mg, 72%).

<sup>1</sup>H NMR Spectrum: (CDCl<sub>3</sub>) 1.98 (m, 2H), 2.08 (m, 2H), 2.83 (d, 3H), 3.32 (m, 2H), 3.72 (m, 2H), 4.01 (s, 3H), 4.48 (m, 1H), 4.64 (m, 1H), 7.16 (m, 2H), 7.23 (s, 1H), 7.31 (s, 1H), 7.38 (br s, 1H), 8.44 (m, 1H), 8.70 (s, 1H); Mass spectrum: MH<sup>+</sup> 460.

**Example 3****Preparation of 4-(3-Chloro-2-fluoroanilino)-7-methoxy-6-[[1-(N-(2-pyrrolidin-1-ylethyl)carbamoyl)piperidin-4-yl]oxy]quinazoline**

A mixture of 6-[[1-(N-(2-chloroethyl)carbamoyl)piperidin-4-yl]oxy}-4-(3-chloro-2-fluoroanilino)-7-methoxyquinazoline (204 mg, 0.4 mmol), pyrrolidine (0.14 ml, 1.6 mmol) and potassium iodide (134 mg, 0.8 mmol) in dimethylacetamide (3 ml) was heated at 80°C for 4 hours. After cooling and evaporation of the solvents under vacuum, the residue was partitioned in water, dichloromethane and extracted with dichloromethane. The organic layer was washed with water and brine, and dried over magnesium sulfate. After evaporation of the solvents under vacuum, the residue was purified by chromatography on silica gel (eluant: 3%

to 4% 7N methanolic ammonia in dichloromethane) to give the title compound as a white solid (77 mg, 36%).

<sup>1</sup>H NMR Spectrum: (CDCl<sub>3</sub>) 1.78 (m, 4H), 1.93 (m, 2H), 2.04 (m, 2H), 2.53 (m, 4H), 2.62 (t, 2H), 3.33 (m, 4H), 3.75 (m, 2H), 4.01 (s, 3H), 4.64 (m, 1H), 5.27 (m, 1H), 7.16 (m, 2H), 7.22 (s, 1H), 7.30 (s, 1H), 7.36 (br s, 1H), 8.45 (m, 1H), 8.70 (s, 1H); Mass spectrum: MH<sup>+</sup> 543.

The 6-([1-(N-(2-chloroethyl)carbamoyl)piperidin-4-yl]oxy)-4-(3-chloro-2-fluoroanilino)-7-methoxyquinazoline used as starting material was made similarly to Example 2 by reaction of 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-  
10 [(piperidin-4-yl)oxy]quinazoline (160 mg, 0.4 mmol) and 2-chloroethylisocyanate (34 µl, 0.4 mmol). Yield: 200 mg, 100%. Mass spectrum: MH<sup>+</sup> 508, 510.

#### Example 4

##### Preparation of 4-(3-Chloro-2-fluoroanilino)-7-methoxy-6-([1-(morpholin-4-ylcarbonyl)piperidin-4-yl]oxy)quinazoline

15 4-Morpholinylcarbonyl chloride (35 µl, 0.3 mmol) was added dropwise to a ice-cooled mixture of 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-[(piperidin-4-yl)oxy]quinazoline (120 mg, 0.3 mmol) and diisopropylethylamine (63 µl, 0.36 mmol) in dichloromethane (5 ml). At the end of the addition, the mixture was stirred at room temperature for 18 hours. The mixture was diluted with dichloromethane, washed with water and brine and dried over magnesium  
20 sulfate. After evaporation of the solvents under vacuum, the residue was purified by chromatography on silica gel (eluant: 1% to 2% 7N methanolic ammonia in dichloromethane) to give the title compound as a white solid (100 mg, 64%).

<sup>1</sup>H NMR Spectrum: (CDCl<sub>3</sub>) 1.93 (m, 2H), 2.05 (m, 2H), 3.20 (m, 2H), 3.29 (m, 4H), 3.62 (m, 2H), 3.70 (m, 4H), 4.01 (s, 3H), 4.64 (m, 1H), 7.16 (m, 2H), 7.20 (s, 1H), 7.31 (m, 2H), 8.49 (m, 1H), 8.71 (s, 1H); Mass spectrum: MH<sup>+</sup> 516.

#### Example 5

##### 4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-([1-(N-methylcarbamoylmethyl)piperidin-4-yl]oxy)quinazoline

2-Chloro-N-methylacetamide (51 mg, 0.47 mmol) was added dropwise to a mixture of  
30 4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-[(piperidin-4-yl)oxy]quinazoline (200 mg, 0.47 mmol), potassium iodide (79 mg, 0.47 mmol) and potassium carbonate (79 mg, 0.57 mmol) in dimethylacetamide (5 ml). The mixture was heated at 70°C for one hour. After cooling and filtration of the solids, the filtrate was purified on an HPLC column (C18, 5 microns, 19 mm

diameter, 100 mm length) of a preparative HPLC-MS system eluting with a mixture of water and acetonitrile containing 2g/l of ammonium formate (gradient) to give the title compound (55 mg, 24%) as a white solid.

**<sup>1</sup>H NMR Spectrum:** (CDCl<sub>3</sub>) 1.98 (m, 2H), 2.07 (m, 2H), 2.44 (m, 2H), 2.86 (m, 2H), 2.87 (d, 3H), 3.06 (s, 2H), 4.01 (s, 3H), 4.48 (m, 1H), 7.07 (m, 1H), 7.15 (m, 1H), 7.20 (s, 1H), 7.30 (m, 2H), 8.32 (m, 1H), 8.66 (s, 1H); **Mass spectrum:** MH<sup>+</sup> 492.

4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-[(piperidin-4-yl)oxy]quinazoline used as starting material was made as follows:

3-Chloro-2,4-difluoroaniline (1.7 g, 10.1 mmol) and 5N hydrogen chloride in isopropanol (2 ml) were added to a suspension of *tert*-butyl 4-[(4-chloro-7-methoxyquinazolin-6-yl)oxy]piperidine-1-carboxylate (4 g, 10.1 mmol, PCT-Int. Appl. WO2003082831, AstraZeneca) in isopropanol (50 ml). The mixture was stirred at 80°C for 3 hours. After evaporation of the solvents, the residue was purified by chromatography on silica gel (eluant: 5-10% 7N methanolic ammonia in dichloromethane) to give 4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-[(piperidin-4-yl)oxy]quinazoline (3.63 g, 85%) as a white solid.

**<sup>1</sup>H NMR Spectrum:** (CDCl<sub>3</sub> + CD<sub>3</sub>CO<sub>2</sub>D): 2.15 (m, 2H), 2.30 (m, 2H), 3.34 (m, 2H), 3.47 (m, 2H), 4.01 (s, 3H), 4.91 (m, 1H), 7.03 (m, 1H), 7.58 (m, 2H), 7.90 (s, 1H), 8.55 (s, 1H); **Mass spectrum:** MH<sup>+</sup> 421.

## 20 Examples 6 to 10

A suspension of [4-({4-(3-chloro-2-fluoroanilino)-7-methoxyquinazolin-6-yl}oxy)piperidin-1-yl]acetic acid dihydrochloride salt (212 mg, 0.4 mmol), 1-hydroxybenzotriazole (66 mg, 0.48 mmol), diisopropylethylamine (0.14 ml, 0.8 mmol), the appropriate amine (0.48 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (92 mg, 0.48 mmol) in dichloromethane (5 ml) was stirred for 2 hours. The mixture was washed with water, 10% aqueous sodium bicarbonate and brine and dried over magnesium sulfate. After evaporation of the solvents, the residue was purified by chromatography on silica gel (eluant: 2-3% 7N methanolic ammonia in dichloromethane) and triturated in acetonitrile to give the title compound.

## 30 Example 6

**4-(3-chloro-2-fluoroanilino)-6-[[1-(N-ethylcarbamoylmethyl)piperidin-4-yl]oxy]-7-methoxyquinazoline**

The amine used was ethylamine.



- 63 -

Yield: 47 mg, 24%; <sup>1</sup>H NMR Spectrum: (CDCl<sub>3</sub>) 1.17 (t, 3H), 1.98 (m, 2H), 2.09 (m, 2H), 2.45 (m, 2H), 2.87 (m, 2H), 3.05 (s, 2H), 3.33 (m, 2H), 4.02 (s, 3H), 4.49 (m, 1H), 7.16 (m, 4H), 7.30 (s, 1H), 7.33 (s br, 1H), 8.48 (m, 1H), 8.71 (s, 1H); Mass spectrum: MH<sup>+</sup> 488.

**Example 7**

5 **4-(3-chloro-2-fluoroanilino)-7-methoxy-6-([1-(N-[2-(pyrrolidin-1-yl)ethyl]carbamoylmethyl)piperidin-4-yl]oxy)quinazoline**

The amine used was 1-(2-aminoethyl)pyrrolidine.

Yield: 53 mg, 24%; <sup>1</sup>H NMR Spectrum: (CDCl<sub>3</sub>) 1.80 (m, 4H), 1.98 (m, 2H), 2.07 (m, 2H), 2.45 (m, 2H), 2.53 (m, 4H), 2.62 (t, 2H), 2.87 (m, 2H), 3.07 (s, 2H), 3.40 (m, 2H), 4.02  
10 (s, 3H), 4.48 (m, 1H), 7.16 (m, 3H), 7.31 (m, 2H), 7.55 (s br 1H), 8.50 (m, 1H), 8.71 (s, 1H); Mass spectrum: MH<sup>+</sup> 557.

**Example 8**

**4-(3-chloro-2-fluoroanilino)-7-methoxy-6-([1-(N-(2-methoxyethyl)carbamoylmethyl)piperidin-4-yl]oxy)quinazoline**

15 The amine used was 2-methoxyethylamine.

Yield: 57 mg, 28%; <sup>1</sup>H NMR Spectrum: (CDCl<sub>3</sub>) 1.98 (m, 2H), 2.09 (m, 2H), 2.45 (m, 2H), 2.87 (m, 2H), 3.07 (s, 2H), 3.38 (s, 3H), 3.48 (s, 4H), 4.02 (s, 3H), 4.49 (m, 1H), 7.16 (m, 3H), 7.31 (m, 2H), 7.48 (s br, 1H), 8.49 (m, 1H), 8.71 (s, 1H); Mass spectrum: MH<sup>+</sup> 518.

**Example 9**

20 **4-(3-chloro-2-fluoroanilino)-6-([1-(N-(2-dimethylaminoethyl)carbamoylmethyl)piperidin-4-yl]oxy)-7-methoxyquinazoline**

The amine used was N,N-dimethylethylenediamine.

Yield 79 mg, 37%; <sup>1</sup>H NMR Spectrum: (CDCl<sub>3</sub>) 1.98 (m, 2H), 2.10 (m, 2H), 2.26 (s, 6H), 2.43 (m, 4H), 2.88 (m, 2H), 3.07 (s, 2H), 3.37 (m, 2H), 3.48 (s br, 1H), 4.03 (s, 3H), 4.49  
25 (m, 1H), 7.16 (m, 3H), 7.31 (m, 2H), 7.51 (s br, 1H), 8.49 (m, 1H), 8.71 (s, 1H); Mass spectrum: MH<sup>+</sup> 531.

**Example 10**

**4-(3-chloro-2-fluoroanilino)-7-methoxy-6-([1-[2-(4-methylpiperazin-1-yl)-2-oxoethyl]piperidin-4-yl]oxy)quinazoline**

30 The amine used was N-methylpiperazine.

Yield: 64 mg, 30%; <sup>1</sup>H NMR Spectrum: (CDCl<sub>3</sub>) 1.96 (m, 2H), 2.11 (m, 2H), 2.32 (s, 3H), 2.40 (m, 6H), 2.87 (m, 2H), 3.24 (s, 2H), 3.65 (m, 4H), 4.02 (s, 3H), 4.47 (m, 1H), 7.16 (m, 3H), 7.30 (m, 1H), 7.33 (s br, 1H), 8.48 (m, 1H), 8.70 (s, 1H); Mass spectrum: MH<sup>+</sup> 543.

**Example 11****4-(3-chloro-2-fluoroanilino)-7-methoxy-6-({1-[2-(piperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy)quinazoline**

The procedure according to Examples 6 to 10 was used except that 1-*tert*-butoxycarbonylpiperazine was used as the amine and that after the aqueous work-up, the residue was stirred for 90 minutes in a 1:1 mixture of dichloromethane-trifluoroacetic acid (3 ml) and then purified by HPLC.

Yield: (150 mg from a 0.56 mmol scale, 51%); <sup>1</sup>H NMR Spectrum: (CDCl<sub>3</sub>) 1.96 (m, 2H), 2.11 (m, 2H), 2.41 (m, 2H), 2.87 (m, 6H), 3.23 (s, 2H), 3.59 (m, 4H), 4.01 (s, 3H), 4.46 (m, 1H), 7.16 (m, 3H), 7.29 (s, 1H), 7.41 (s br, 1H), 8.45 (m, 1H), 8.70 (s, 1H); Mass spectrum: MH<sup>+</sup> 529.

The [4-({4-(3-chloro-2-fluoroanilino)-7-methoxyquinazolin-6-yl}oxy)piperidin-1-yl]acetic acid dihydrochloride salt used as starting material was made as follows:

*Tert*-butyl chloroacetate (1.43 ml, 10 mmol) was added dropwise to a mixture of 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-[(piperidin-4-yl)oxy]quinazoline (4.02 g, 10 mmol), potassium iodide (1.66 g, 10 mmol) and potassium carbonate (1.66 g, 12 mmol) in dimethylacetamide (50 ml). The mixture was heated at 70°C for one hour. After evaporation of the solvents under vacuum, the residue was triturated in water. The resulting solid was filtered, washed with water and purified by chromatography on silica gel (eluant: 2% 7N methanolic ammonia in dichloromethane) to give *tert*-butyl [4-({4-(3-chloro-2-fluoroanilino)-7-methoxyquinazolin-6-yl}oxy)piperidin-1-yl]acetate as a white solid (3.0 g, 60%).

NMR Spectrum: (CDCl<sub>3</sub>) 1.48 (s, 9H), 2.01 (m, 2H), 2.10 (m, 2H), 2.56 (m, 2H), 2.89 (m, 2H), 3.19 (s, 2H), 4.01 (s, 3H), 4.49 (m, 1H), 7.16 (m, 3H), 7.29 (m, 2H), 8.48 (m, 1H), 8.70 (s, 1H); Mass spectrum: MH<sup>+</sup> 517.

A suspension of *tert*-butyl [4-({4-(3-chloro-2-fluoroanilino)-7-methoxyquinazolin-6-yl}oxy)piperidin-1-yl]acetate (3.0 g, 5.8 mmol) in a solution of 4N hydrogen chloride in dioxane (40 ml) was stirred at room temperature for 3 hours. The solvents were evaporated under high vacuum. The residue was triturated in ether, filtered and washed with ether to give [4-({4-(3-chloro-2-fluoroanilino)-7-methoxyquinazolin-6-yl}oxy)piperidin-1-yl]acetic acid as the dihydrochloride salt (3.1 g, 100%). Mass spectrum: MH<sup>+</sup> 461.

**Example 12****4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-({1-[2-(4-methylpiperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy)quinazoline**

- 65 -

[4-({4-(3-chloro-2,4-difluoroanilino)-7-methoxyquinazolin-6-yl}oxy)piperidin-1-yl]acetic acid dihydrochloride salt and N-methylpiperazine were converted to the title compound (126 mg, 56%) using the procedure according to Examples 6 to 10.

<sup>1</sup>H NMR Spectrum: (CDCl<sub>3</sub>) 1.94 (m, 2H), 2.09 (m, 2H), 2.31 (s, 3H), 2.40 (m, 6H), 2.84 (m, 2H), 3.23 (s, 2H), 3.65 (m, 4H), 4.01 (s, 3H), 4.45 (m, 1H), 7.06 (m, 1H), 7.22 (s, 1H), 7.29 (m, 1H), 7.36 (s br, 1H), 8.28 (m, 1H), 8.65 (s, 1H); Mass spectrum: MH<sup>+</sup> 561

The [4-({4-(3-chloro-2,4-difluoroanilino)-7-methoxyquinazolin-6-yl}oxy)piperidin-1-yl]acetic acid dihydrochloride salt used as starting material was made from 4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-[(piperidin-4-yl)oxy]quinazoline using the same procedure as described in Example 11:

*tert*-Butyl [4-({4-(3-chloro-2,4-difluoroanilino)-7-methoxyquinazolin-6-yl}oxy)piperidin-1-yl]acetate (2.56 g, 67%): Mass spectrum: MH<sup>+</sup> 535.

[4-({4-(3-chloro-2,4-difluoroanilino)-7-methoxyquinazolin-6-yl}oxy)piperidin-1-yl]acetic acid (dihydrochloride salt, 2.45 g, 93%): Mass spectrum: MH<sup>+</sup> 479.

### 15 Example 13

#### Pharmaceutical Compositions

The following illustrates a representative pharmaceutical dosage forms of the invention as defined herein (the active ingredient being termed "Compound X"), for therapeutic or prophylactic use in humans:

20

(a)	Tablet I	mg/tablet
	Compound X.....	100
	Lactose Ph.Eur.....	182.75
	Croscarmellose sodium.....	12.0
25	Maize starch paste (5% w/v paste).....	2.25
	Magnesium stearate.....	3.0

30


(b)	Injection I	(50 mg/ml)
	Compound X.....	5.0% w/v
	1M Sodium hydroxide solution.....	15.0% v/v
	0.1M Hydrochloric acid (to adjust pH to 7.6)	
	Polyethylene glycol 400.....	4.5% w/v
	Water for injection to 100%.	

- 66 -

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. For example the tablet may be prepared by blending the components together and compressing the mixture into a tablet.

**CLAIMS**

1. A quinazoline derivative of the Formula I:



I

- 5 wherein  $n$  is 0, 1, 2 or 3,  
 each  $R^5$  is independently selected from halogeno, cyano, nitro, hydroxy, amino, carboxy, sulfamoyl, trifluoromethyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl,  
 10  $\underline{N}$ -(1-6C)alkylsulfamoyl, and  $\underline{N,N}$ -di-[(1-6C)alkyl]sulfamoyl,  $C(O)NR^6R^7$  where  $R^6$  and  $R^7$  are independently selected from hydrogen, optionally substituted (1-6C)alkyl, optionally substituted (3-8C)cycloalkyl or optionally substituted aryl, or  $R^6$  and  $R^7$  together with the nitrogen to which they are attached form an optionally substituted heterocyclic ring which may contain additional heteroatoms;  
 15  $X^1$  is a direct bond or O;  
 $R^1$  is selected from hydrogen and (1-6C)alkyl, wherein the (1-6C)alkyl group is optionally substituted by one or more substituents, which may be the same or different, selected from hydroxy and halogeno, and/or a substituent selected from amino, nitro, carboxy, cyano, halogeno, (1-6C)alkoxy, hydroxy(1-6C)alkoxy, (2-8C)alkenyl, (2-8C)alkynyl,  
 20 (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, carbamoyl,  $\underline{N}$ -(1-6C)alkylcarbamoyl,  $\underline{N,N}$  di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino,  $\underline{N}$ -(1-6C)alkyl-(2-6C)alkanoylamino, (1-6C)alkoxycarbonyl, sulfamoyl,  $\underline{N}$ -(1-6C)alkylsulfamoyl,  $\underline{N,N}$ -di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and  
 25  $\underline{N}$ -(1-6C)alkyl-(1-6C)alkanesulfonylamino;  
 $m$  is 0, 1, 2 or 3;

- 68 -

$R^2$  is hydrogen or (1-6C)alkyl; and

$R^3$  is (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl or (1-6C)alkoxy, any of which can be optionally substituted on a carbon atom by a (1-6C)alkoxy, amino, (1-6C)alkylamino, di-(1-6C)alkylamino, or a group  $S(O)_s(1-6C)alkyl$  where  $s$  is 0, 1 or 2, or a saturated 5 or 6 membered heterocyclic ring which optionally contains additional heteroatoms selected from oxygen, sulfur or  $NR^8$  where  $R^8$  is hydrogen, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkylsulfonyl or (1-6C)alkylcarbonyl; or  $R^2$  and  $R^3$  together with the nitrogen atom to which they are attached form a saturated 5 or 6 membered heterocyclic ring which optionally contains additional heteroatoms selected from oxygen, S, SO or  $S(O)_2$  or  $NR^8$  where  $R^8$  is as defined above; provided that the quinazoline derivative is not:

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy}-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonyl]-piperidin-4-yl-oxy}-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(diethylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(piperidin-1-yl)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(pyrrolidin-1-yl)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(4-methyl-piperazin-1-yl)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy}-7-ethoxy-quinazoline;

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy}-7-(2-methoxy-ethoxy)-quinazoline;


4-[(3-ethynyl-phenyl)amino]-6-{1-[(morpholin-4-yl)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;

- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(ethylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(isopropylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 5 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonylmethyl]-piperidin-4-yl-oxy}-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonylmethyl]-piperidin-4-yl-oxy}-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 10 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(methylamino)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 15 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(dimethylamino)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(pyrrolidin-1-yl)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonylmethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 20 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(methylamino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(2-methoxyethyl)amino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 25 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(N-methyl-N-2-methoxyethyl)amino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(3-methoxypropyl)amino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(N-methyl-N-3-methoxypropyl)amino)carbonyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;
- 30 4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonylethyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline; or

- 70 -

4-[(3-chloro-4-fluorophenyl)amino]-6-{1-[(morpholin-4-yl)carbonylpropyl]-piperidin-4-yl-oxy}-7-methoxy-quinazoline;  
or a pharmaceutically acceptable salt thereof.


- 5 2. A quinazoline derivative according to claim 1, wherein n is 1, 2 or 3.
3. A quinazoline derivative according to claim 1 or claim 2, wherein n is 2 or 3.
4. A quinazoline derivative according to any one of claims 1 to 3, wherein n is 2.
- 10 5. A quinazoline derivative according to any one of claims 1 to 3, wherein n is 3.
6. A quinazoline derivative according to any one of the preceding claims, wherein each group  $R^5$  is a halogeno group.
- 15 7. A quinazoline derivative according to any one of the preceding claims, wherein each group  $R^5$  is selected from chloro and fluoro.
8. A quinazoline derivative according to any one of the preceding claims, which includes
- 20 a group  $R^5$  positioned at an ortho- (2-) position on the benzene ring to which it is attached.
9. A quinazoline derivative according to claim 8, wherein the group  $R^5$  positioned at the ortho- (2-) position is fluoro.
- 25 10. A quinazoline derivative according to any one of the preceding claims, wherein in the Formula I, the group of sub-formula (i):



is a group of sub-formula (ii):



- 71 -



(ii)

wherein (a) one of  $R^{10}$  or  $R^{12}$  is hydrogen and the other is halogeno, and  $R^{11}$  is halogeno, or (b)  $R^{10}$  is halogeno,  $R^{11}$  is halogeno and  $R^{12}$  is selected from hydrogen or halogeno, or (c)  $R^{10}$  is fluoro,  $R^{11}$  is chloro, and  $R^{12}$  is selected hydrogen or fluoro.

5

11. A quinazoline derivative according to claim 10, wherein one of  $R^{10}$  or  $R^{12}$  is hydrogen and the other is fluoro, and  $R^{11}$  is chloro.

12. A quinazoline derivative according to claim 10, wherein  $R^{10}$  is fluoro,  $R^{11}$  is chloro,  
10 and  $R^{12}$  is hydrogen.

13. A quinazoline derivative according to claim 10, wherein  $R^{10}$  is fluoro,  $R^{11}$  is chloro, and  $R^{12}$  is fluoro.

15 14. A quinazoline derivative according to any one of the preceding claims, wherein  $X^1$  is oxygen.

15. A quinazoline derivative according to any one of the preceding claims, wherein  $R^1$  is selected from hydrogen, (1-6C)alkyl and (1-6C)alkoxy(1-6C)alkyl, wherein any (1-6C)alkyl  
20 group in  $R^1$  optionally bears one or more hydroxy or halogeno substituents


16. A quinazoline derivative according to claim 15, wherein  $R^1$  is selected from (1-6C)alkyl, which optionally bears one or more hydroxy or halogeno substituents.

25 17. A quinazoline derivative according to any one of the claims 1 to 13, wherein  $R^1-X^1$  is selected from hydrogen, methoxy, ethoxy and 2-methoxyethoxy.

18. A quinazoline derivative according to claim 17, wherein  $R^1-X^1$  is methoxy.

- 72 -

19. A quinazoline derivative according to claim 1 of Formula IA:




IA

wherein  $R^2$ ,  $R^3$  and  $m$  are as defined in claim 1,  $R^{10}$ ,  $R^{11}$  and  $R^{12}$  are as defined in any one of claims 10 to 13, and  $R^{13}$  is selected from hydrogen, methoxy, ethoxy and 2-methoxyethoxy.

5

20. A quinazoline derivative according to claim 1 of Formula IB:




IB

wherein  $R^2$ ,  $R^3$  and  $m$  are as defined in claim 1 and  $R^{13}$  is selected from hydrogen, methoxy, ethoxy and 2-methoxyethoxy.

10

21. A quinazoline derivative according to claim 1 of Formula IC:



IC

- 73 -

wherein  $R^2$ ,  $R^3$  and  $m$  are as defined in claim 1 and  $R^{13}$  is selected from hydrogen, methoxy, ethoxy and 2-methoxyethoxy.

22. A quinazoline derivative according to any one of claims 19 to 21, wherein  $R^{13}$  is  
5 methoxy.
23. A quinazoline derivative according to any one of the preceding claims, wherein  $m$  is 0 or 1.
- 10 24. A quinazoline derivative according to any one of the preceding claims, wherein  $m$  is 1.
25. A quinazoline derivative according to any one of the preceding claims, wherein  $R^2$  is hydrogen or (1-3C)alkyl.
- 15 26. A quinazoline derivative according to any one of the preceding claims, wherein  $R^2$  is hydrogen or methyl.
27. A quinazoline derivative according to any one of the preceding claims, wherein  $R^2$  is  
20 hydrogen.
28. A quinazoline derivative according to any one of the preceding claims, wherein  $R^3$  is (1-6C)alkyl.
29. A quinazoline derivative according to any one of the preceding claims, wherein  $R^3$  is  
25 (1-3C)alkyl.
30. A quinazoline derivative according to any one of the preceding claims, wherein  $R^3$  is methyl.
- 30 31. A quinazoline derivative according to claim 1, which is selected from one or more of the following:  
4-(3-chloro-2-fluoroanilino)-7-methoxy-6- $\{[1-(N\text{-methylcarbamoylmethyl})\text{piperidin-4-yl}]\text{-oxy}\}$ quinazoline;


- 74 -

- 4-(3-chloro-2-fluoroanilino)-6-{{1-(N,N-dimethylcarbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(morpholin-4-ylcarbonylmethyl)piperidin-4-yl}oxy}-quinazoline;
- 5 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(pyrrolidin-1-ylcarbonyl)piperidin-4-yl}oxy}quinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoyl)piperidin-4-yl}oxy}quinazoline;
- 4-(3-chloro-2-fluoroanilino)-6-{{1-(N-(2-dimethylaminoethyl)carbamoyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;
- 10 4-(3-chloro-2-fluoroanilino)-6-{{1-(N,N-dimethylcarbamoyl)piperidin-4-yl}oxy}7-methoxyquinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(morpholin-4-ylcarbonyl)piperidin-4-yl}oxy}quinazoline;
- 15 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-[2-pyrrolidin-1-ylethyl]carbamoyl)piperidin-4-yl}oxy}quinazoline;
- 4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-{{1-(N-methylcarbamoylmethyl)piperidin-4-yl}oxy}quinazoline;
- 4-(3-chloro-2-fluoroanilino)-6-{{1-(N-ethylcarbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;
- 20 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-[2-(pyrrolidin-1-yl)ethyl]carbamoylmethyl)piperidin-4-yl}oxy}quinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-(N-(2-methoxyethyl)carbamoylmethyl)piperidin-4-yl}oxy}quinazoline;
- 25 4-(3-chloro-2-fluoroanilino)-6-{{1-(N-(2-dimethylaminoethyl)carbamoylmethyl)piperidin-4-yl}oxy}-7-methoxyquinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-[2-(4-methylpiperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy}quinazoline;
- 4-(3-chloro-2-fluoroanilino)-7-methoxy-6-{{1-[2-(piperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy}quinazoline; and
- 30 4-(3-chloro-2,4-difluoroanilino)-7-methoxy-6-{{1-[2-(4-methylpiperazin-1-yl)-2-oxoethyl]piperidin-4-yl}oxy}quinazoline;
- or a pharmaceutically acceptable salt thereof.

- 75 -


32. A process for preparing a quinazoline derivative according to any one of the preceding claims, which comprises either

**Process (a)** reacting a compound of the Formula II:



II

5 wherein R¹, X¹, R⁵ and n have any of the meanings defined in claim 1 except that any functional group is protected if necessary,  
with a compound of the Formula III:




III

wherein R², R³ and m have any of the meanings defined in claim 1 except that any functional  
10 group is protected if necessary and Lg is a displaceable group, wherein the reaction is conveniently performed in the presence of a suitable base,

**Process (b)** modifying a substituent in or introducing a substituent into another quinazoline derivative of Formula I or a pharmaceutically acceptable salt thereof, as hereinbefore defined except that any functional group is protected if necessary;



15 **Process (c)** reacting a compound of Formula IV:

- 76 -



IV

where  $R^1$ ,  $X^1$ ,  $R^5$  and  $n$  are as defined in relation to Formula I except that any functional group is protected if necessary, with a compound of the Formula V or V':



5. wherein  $R^2$  and  $R^3$  are as defined above and  $m'$  is 0, 1, 2 or 3, provided that it is not 0 when  $R^2$  is hydrogen, and Lg is a displaceable group;

**Process (d)** removal of a protecting group from a quinazoline derivative of Formula I, or a pharmaceutically acceptable salt thereof;


**Process (e)** reacting a compound of the Formula II as hereinbefore defined with a

- 10 compound of the Formula III as defined hereinbefore except Lg is OH under Mitsunobu conditions;

**Process (f)** for the preparation of those compounds of the Formula I wherein  $R^1-X^1$  is a hydroxy group, cleavage of a quinazoline derivative of the Formula I wherein  $R^1-X^1$  is a (1-6C)alkoxy group;

- 15 **Process (g)** for the preparation of those compounds of the Formula I wherein  $X^1$  is O and  $R^1$  is not hydrogen, by the reaction of a compound of the Formula VI:

- 77 -




VI

wherein  $R^2$ ,  $R^3$ ,  $R^5$ ,  $m$  and  $n$  have any of the meanings defined in claim 1 except that any functional group is protected if necessary, with a compound of the formula  $R^1$ -Lg, wherein  $R^1$  has any of the meanings defined in claim 1 except that it is not hydrogen and except that any

5 functional group is protected if necessary and Lg is a displaceable group;

**Process (h)** for the preparation of those compounds of the Formula I wherein  $R^1$  contains a (1-6C)alkoxy or substituted (1-6C)alkoxy group or a (1-6C)alkylamino or substituted (1-6C)alkylamino group, alkylation of a quinazoline derivative of the Formula I wherein or  $R^1$  contains a hydroxy group or a primary or secondary amino group as appropriate;

10 **Process (i)** for the preparation of those compounds of the Formula I wherein  $R^1$  is substituted by a group T, wherein T is selected from (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (2-6C)alkanoylamino, (1-6C)alkylthio, (1-6C)alkylsulfinyl and (1-6C)alkylsulfonyl, the reaction of a compound of the Formula VII:




VII

15 wherein  $R^2$ ,  $R^3$ ,  $R^5$ ,  $X^1$ ,  $n$  and  $m$  have any of the meanings defined hereinbefore except that any functional group is protected if necessary,  $R^{1'}$  is a group  $R^1$  as defined herein except that any T groups are replaced with Lg, and Lg is a displaceable group (for example chloro or bromo) with a compound of the formula TH, wherein T is as defined above except that any functional group is protected if necessary;


- 78 -

Process (j) reacting a compound of the Formula VIII:



VIII


wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $X^1$ , and  $m$  have any of the meanings defined in claim 1 except that any functional group is protected if necessary and  $Lg$  is a displacable group as hereinbefore  
 5 defined,  
 with an aniline of the Formula IX:



IX

wherein  $R^5$  and  $n$  have any of the meanings defined in claim 1 except that any functional group is protected if necessary, and wherein the reaction is conveniently performed in the  
 10 presence of a suitable acid;

Process (k) for the preparation of those compounds of the Formula I wherein  $m$  is 1, 2 or 3, coupling of a compound of Formula X:




X

wherein  $m$  is 1, 2 or 3 and  $R^1$ ,  $X^1$ ,  $R^5$ , and  $n$  are as hereinbefore defined in claim 1, except  
 15 any functional group is protected if necessary, with a primary or secondary amine of formula  $R^2NHR^3$  where  $R^2$  and  $R^3$  are as defined in claim 1;




- 79 -

**Process (l)** By reacting a compound of Formula IV as defined above except that any functional group is protected if necessary, with a compound of the Formula V'':



using a reductive amination procedure,

- 5 **Process (m)** for the preparation of those compounds of the Formula I wherein R<sup>3</sup> is (2-6C)alkyl substituted on a carbon atom by an amino, (1-6C)alkylamino, di-(1-6C)alkylamino or a saturated 5 or 6 membered heterocyclic ring which contains NR<sup>8</sup> where R<sup>8</sup> is as defined in claim 1, by reacting a compound of the Formula XX:



- 10 wherein R<sup>3a</sup> is Lg-(2-6C)alkyl, wherein Lg is a displaceable group and wherein R<sup>1</sup>, R<sup>2</sup>, X<sup>1</sup>, R<sup>5</sup>, m and n have any of the meanings defined hereinbefore except that any functional group is protected if necessary,

with ammonia or with a suitable primary or secondary amine,

and whereafter any of said processes, any protecting group that is present is removed.

15

33. A pharmaceutical composition which comprises a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined in any one of claims 1 to 31 in association with a pharmaceutically-acceptable diluent or carrier.

- 20 34. A quinazoline derivative of the Formula I as defined in any one of claims 1 to 31, or a pharmaceutically acceptable salt thereof, for use as a medicament.

- 80 -

35. The use of a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof, as defined in any one of claims 1 to 31 in the manufacture of a medicament for use in the production of an anti-proliferative effect in a warm-blooded animal.
- 5 36. A method for producing an anti-proliferative effect in a warm-blooded animal in need of such treatment which comprises administering to said animal a quinazoline derivative of the Formula I, or a pharmaceutically acceptable salt thereof, as defined any one of claims 1 to 31.
- 10 37. A compound of the Formula VI, VII, VIII, X or XX as defined in claim 32 or a salt thereof.

# INTERNATIONAL SEARCH REPORT

GB2004/003937

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/12 A61K31/517 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, EMBASE, BIOSIS, BEILSTEIN Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 02/41882 A (NOVARTIS ERFIND VERWALT GMBH ; NOVARTIS AG (CH); TRAXLER PETER (CH); B) 30 May 2002 (2002-05-30) cited in the application page 17; claim 8	1-37
A	EP 0 566 226 A (ZENECA LTD) 20 October 1993 (1993-10-20) cited in the application page 10, line 11 - line 20; claim 1	1-37
	----- -/-- -----	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

17 January 2005

Date of mailing of the international search report

25/01/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Härtinger, S

## INTERNATIONAL SEARCH REPORT

GB2004/003937

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	HENNEQUIN L F ET AL: "Novel 4-anilinoquinazolines with C-7 basic side chains: Design and structure activity relationship of a series of potent, orally active, VEGF receptor tyrosine kinase inhibitors" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 45, no. 6, 14 March 2002 (2002-03-14), pages 1300-1312, XP002256124 ISSN: 0022-2623 figure 2	1-37
P,X	----- WO 03/082290 A (SOLCA FLAVIO ; BOEHRINGER INGELHEIM PHARMA (DE); HIMMELSBACH FRANK (DE) 9 October 2003 (2003-10-09) cited in the application pages 58,89,91; claims 3-5 page 100 - page 102	1-37
P,X	----- WO 03/082831 A (ASTRAZENECA UK LTD ; BRADBURY ROBERT HUGH (GB); HENNEQUIN LAURENT FRAN) 9 October 2003 (2003-10-09) cited in the application page 162 - page 164; claim 7 -----	1-37

# INTERNATIONAL SEARCH REPORT

Information on patent family members

GB2004/003937

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0241882	A	30-05-2002	AU 2368402 A	03-06-2002
			CA 2427184 A1	30-05-2002
			WO 0241882 A2	30-05-2002
			EP 1339458 A2	03-09-2003
			JP 2004513964 T	13-05-2004
			US 2004034026 A1	19-02-2004
EP 0566226	A	20-10-1993	AT 130000 T	15-11-1995
			AU 661533 B2	27-07-1995
			AU 3101093 A	22-07-1993
			CA 2086968 A1	21-07-1993
			CZ 9300043 A3	13-10-1993
			DE 69300754 D1	14-12-1995
			DE 69300754 T2	28-03-1996
			DK 566226 T3	18-03-1996
			EP 0566226 A1	20-10-1993
			ES 2078798 T3	16-12-1995
			FI 930208 A	21-07-1993
			GR 3018143 T3	29-02-1996
			HK 36497 A	04-04-1997
			HU 63153 A2	28-07-1993
			HU 9500185 A3	28-07-1995
			IL 104479 A	22-12-1999
			KR 229294 B1	01-11-1999
			MX 9300277 A1	30-06-1994
			NO 930178 A	21-07-1993
			NZ 245662 A	26-09-1995
			RU 2127263 C1	10-03-1999
			SK 1693 A3	09-09-1993
			US 5457105 A	10-10-1995
			US 5616582 A	01-04-1997
			ZA 9300015 A	20-07-1993
			JP 2994165 B2	27-12-1999
			JP 6073025 A	15-03-1994
WO 03082290	A	09-10-2003	DE 10214412 A1	09-10-2003
			DE 10231711 A1	22-01-2004
			CA 2476008 A1	09-10-2003
			WO 03082290 A1	09-10-2003
			EP 1492536 A1	05-01-2005
			US 2004048880 A1	11-03-2004
WO 03082831	A	09-10-2003	CA 2479642 A1	09-10-2003
			EP 1487806 A1	22-12-2004
			WO 03082831 A1	09-10-2003